



TECHNICAL DOCUMENT

European surveillance of *Clostridium difficile* infections

Surveillance protocol version 2.1

ECDC TECHNICAL DOCUMENT

European surveillance of *Clostridium difficile* infections

Surveillance protocol version 2.1



This technical document was commissioned by the European Centre for Disease Prevention and Control (ECDC), coordinated by Pete Kinross and Carl Suetens, and produced by the ECDIS-Net project coordinated by the Leiden University Medical Center LUMC, The Netherlands.

Contributing authors

Axel Kola, Michael Behnke, Petra Gastmeier (Charité - Universitätsmedizin Berlin, Germany); Sofie M van Dorp, Ed J Kuijper (Leiden University Medical Center, The Netherlands); Pete Kinross, Carl Suetens (ECDC); ECDIS-Net surveillance working group (see Acknowledgements).

In accordance with the Staff Regulations for Officials and Conditions of Employment of Other Servants of the European Union and the ECDC Independence Policy, ECDC staff members shall not, in the performance of their duties, deal with a matter in which, directly or indirectly, they have any personal interest such as to impair their independence.

Declarations of interest were received from the contractors in accordance with ECDC's Independence Policy and no conflict was identified.

This technical document was sent for consultation to the ECDC National Focal Points for Healthcare-Associated Infections and to the ECDC Advisory Forum.

Acknowledgements

ECDC would like to thank the other members of the ECDIS-Net surveillance working group for contributing to the CDI protocol meetings (Berlin, 27 Feb 2012; Berlin (ECCMID), 27 Apr 2013; Leiden, 22 Jan 2014), organising the pilot surveillance study in 2013 and/or commenting on the document: D Schmid, E Simons, F Allerberger (Austria); J van Broeck, M Delmée (Belgium); V Jindrák (Czech Republic); K EP Olsen (Denmark), A Pavelkovich, M Altmets (Estonia), O Lyytikäinen, S Mentula (Finland), F Barbut, A Collignon, B Coignard, S Vaux, K Chami (France), D Weitzel-Kage (Germany), A Hajdu, A Kurcz, K Borocz, K Antmann, Zs Barna (Hungary), A Ingebretsen, E Lingaas (Norway), D Notermans, S C de Greef, B van Benthem (the Netherlands), H Pituch, P Karpinski (Poland), I S Macovei (Romania), M Drakulovic, M Jovanovic (Serbia), T Åkerlund, J Struwe (Sweden), C Wiuff, J Coia, T Morris, M Wilcox (United Kingdom), Hospital Contact Points of the pilot CDI surveillance study.

Suggested citation: European Centre for Disease Prevention and Control. European Surveillance of *Clostridium difficile* infections. Surveillance protocol version 2.1 Stockholm: ECDC; 2015.

Stockholm, May 2015

ISBN 978-92-9193-635-9

doi 10.2900/071325

Catalogue number TQ-04-15-291-EN-N

© European Centre for Disease Prevention and Control, 2015

Reproduction is authorised, provided the source is acknowledged

Contents

Abbreviations	iv
Objectives	1
Objectives of CDI surveillance in the EU.....	1
Objectives of this protocol	1
Definitions and inclusion/exclusion criteria.....	2
Hospitals.....	2
Wards.....	2
Long-term care facility	2
Patient (denominator) data	2
Definition of <i>Clostridium difficile</i> infection (CDI).....	2
Case (numerator) data	3
Recurrent CDI cases	3
CDI case origin	3
Data collection: the three options	4
Who collects the data?	4
Form H: Hospital-based data.....	5
Definitions.....	5
Form E: Additional case-based data	10
Definitions.....	10
Definitions.....	11
References	13
Annex - Ward/unit specialty code list	14

Abbreviations

APACHE	Acute Physiology and Chronic Health Evaluation
ARHAI	Antimicrobial resistance and healthcare-associated infections
CA	Clinically-associated
CDI	<i>Clostridium difficile</i> infection
ECDIS-Net	European <i>Clostridium difficile</i> Infection Network project
EIA	Enzyme immunoassay
HA	Hospital-associated
ICU	Intensive care unit
MIC	Minimum inhibitory concentration
NAAT	Nucleic acid amplification test
PCR	Polymerase chain reaction
TcdA	<i>Clostridium difficile</i> toxin A
TcdB	<i>Clostridium difficile</i> toxin B
UNK	Unknown

Background

In response to the emerging problems with *Clostridium difficile* infections (CDIs), the European Centre for Disease Prevention and Control (ECDC) in collaboration with the US Centres for Disease Control and Prevention (CDC), published background information about the changing epidemiology of CDIs, agreed on CDI case-definitions and issued recommendations for the surveillance of CDIs [1]. An ECDC-funded survey performed in 2008 [2] revealed a mean incidence of 4.1 per 10 000 patient-days per hospital (range: 0.0–36.3), almost 70% higher than that reported in a previous European surveillance study [3] performed in 2005 (2.45 per 10 000 patient-days per hospital, range: 0.13–7.1), although the surveys each had a different design. Standardised periodic or continuous surveillance of the incidence of CDI is more likely to facilitate the identification of epidemiological changes and is an essential tool for CDI prevention and control. Microbiological data may be an important supplement to surveillance data and allow further insights into epidemiological changes. However, molecular typing and antimicrobial susceptibility testing of isolates are mainly restricted to outbreaks of *C. difficile* or severe cases of CDI.

Facing the lack of standardised surveillance of CDI in EU Member States, ECDC launched a call for tender to support capacity building for surveillance of *Clostridium difficile* infections at the European level in 2010. The contract was awarded to a consortium that carried out the European *Clostridium difficile* Surveillance Network (ECDIS-Net) project [4]¹. The ECDIS-Net project developed a protocol for the surveillance of CDI, composed of three options:

- a 'minimal' surveillance option to collect aggregated denominator and numerator data at the hospital level
- a 'light' surveillance option to collect additional case-based data on CDI cases in hospitals
- an 'enhanced' surveillance option to collect some additional case-based data characteristics and detailed microbiological data (typing and susceptibility testing) on a maximum of 10 patients per surveillance period.

Objectives

Objectives of CDI surveillance in the EU

The objectives for the surveillance of CDIs are:

- to estimate the incidence of CDIs in European acute care hospitals
- to assess the burden of CDIs (including recurrent CDI cases) in European acute care hospitals
- to provide participating hospitals with a standardised tool to measure and compare their own incidence rates with those observed in other participating hospitals
- to assess adverse outcomes of CDIs including death
- to describe the epidemiology of *C. difficile* at the local, national and European level, in terms of factors such as antibiotic susceptibility, PCR ribotype, presence of *Clostridium difficile* toxin A (TcdA), *Clostridium difficile* toxin B (TcdB) and binary toxin, morbidity and mortality of infection, and the detection of new/emerging types.

Objectives of this protocol

This protocol prescribes the methodology, and provides the data collection tools required to achieve the objectives of European surveillance of CDIs. This requires national or regional coordinators to choose one of three CDI surveillance options for data collection by surveyors at the hospital level. Each option corresponds to the collection of progressively more detailed information:

- the minimal CDI surveillance option corresponds to collection of only aggregated numerator and denominator data
- the light surveillance option necessitates collection of case-based numerator data
- the enhanced surveillance option necessitates collection of additional case-based data as well as microbiological data, i.e. molecular characterisation and antimicrobial susceptibility testing data, for the isolates corresponding to the first 10 consecutively detected CDI cases in each healthcare facility (see section 'Data collection').

¹ Consortium composed of Leiden University Medical Center, The Netherlands (E.J. Kuijper, coordination), University of Leeds & Health Protection Agency, England, United Kingdom (M. Wilcox), University Hospital of Wales, Cardiff, United Kingdom (V. Hall), Centre for Infectious Disease Control, RIVM, Bilthoven, The Netherlands (D. Notermans), Charité - Universitätsmedizin Berlin, Germany (P. Gastmeier, A. Kola), in collaboration with ECDC (C. Suetens, K. Weist, P. Kinross).

Definitions and inclusion/exclusion criteria

This section provides definitions and inclusion/exclusion criteria for reference. It is recommended that they are read before surveillance activities. The definition of each variable collected using a surveillance form is provided within the section of this protocol dedicated to that particular form.

Hospitals

An acute care hospital is defined according to national definitions. All acute care hospitals are eligible for inclusion. There is no minimum size of hospitals.

It is preferable for hospitals with more than one geographical site to report each site that has a separate infection control team/unit separately, if this is practicable. Otherwise, it is sufficient to report for the entire hospital group.

The participation of hospitals to the national surveillance of CDI may be voluntary or mandatory, depending on the country. Representative sampling of hospitals is not required but is recommended.

Wards

Include all wards in acute care facilities, including long-term care wards. Exclusion of wards is not allowed.

Long-term care facility

A long-term care facility is defined as a facility in which residents need constant supervision (24 hours); need 'high-skilled nursing care' (i.e. more than 'basic' nursing care and assistance for daily living); are medically stable and do not need constant 'specialised medical care' (i.e. administered by specialised physicians); and do not need invasive medical procedures (e.g. ventilation).

Patient (denominator) data

All hospitalised patients should be included in the denominator, including children age two years or less. A patient is considered as hospitalised when he or she is registered as such in the local hospital administration system and will therefore contribute to the denominator data (number of admissions or discharges, number of patient-days). Usually, this involves at least one overnight stay in the hospital.

Definition of *Clostridium difficile* infection (CDI)

A case of *Clostridium difficile* infection (CDI) (previously also referred to as *C. difficile* associated diarrhoea or CDAD) must meet at least one of the following criteria [1]:

- diarrhoeal stools or toxic megacolon AND a positive laboratory assay for *C. difficile* toxin A and/or B in stools or a toxin-producing *C. difficile* organism detected in stool via culture or other means e.g. a positive PCR result;
OR
- pseudomembranous colitis revealed by lower gastro-intestinal endoscopy;
OR
- colonic histopathology characteristic of *C. difficile* infection (with or without diarrhoea) on a specimen obtained during endoscopy, colectomy or autopsy.

Case (numerator) data

Numerator data are collected for all hospitalised patients that meet the definition of CDI, and meet at least one of the following inclusion criteria.

Inclusion criteria:

- the date of CDI symptom onset was within the surveillance period (even if the patient was admitted before the start of the surveillance period)
OR
- the patient was admitted to the hospital during the surveillance period with signs and symptoms of CDI present at admission, even if this episode of CDI was already diagnosed prior to admission (e.g. at the outpatient department)
OR
- recurrent cases of CDI (see definition below).

Exclusion criteria:

- Day cases, e.g. one day surgery; patients in the emergency room; dialysis patients (outpatients).

It is recognised that many children are asymptomatically colonised with *C. difficile*. Detection of *C. difficile* in children of less than two years of age should only lead to the inclusion of these patients as CDI cases in the numerator if there is compelling clinical evidence for CDI.

Recurrent CDI cases

In clinical practice, it is not possible to differentiate between a relapse involving the same strain and re-infection with a different strain. The term 'recurrence' is used as a designation for both.


Recurrent CDI cases are patients meeting the CDI case definition with an episode of CDI (return of diarrhoeal stools with a positive laboratory test after the end of treatment) more than two weeks and less than eight weeks following the onset of a previous episode.

CDI cases with symptom onset more than eight weeks after the onset of a previous episode are included as new CDI cases.

CDI case origin

The origin of a CDI case can be healthcare-associated, community-associated or unknown.

Figure 1. Designation of CDI cases as healthcare-associated or community-associated based on location and time of onset of symptoms.



Healthcare-associated CDI (HA CDI) is defined as a case of CDI with onset of symptoms:

- on day three or later, following admission to a healthcare facility on day one,
OR
- in the community within four weeks of discharge from a healthcare facility (including the current hospital or a previous stay in any other healthcare facility).

Community-associated CDI (CA CDI) is defined as a case of CDI with onset of symptoms:

- outside of healthcare facilities
AND without discharge from a healthcare facility within the previous 12 weeks,
OR
- on the day of admission to a healthcare facility or on the following two days
AND not resident in a healthcare facility within the previous 12 weeks.

Unknown association: the CDI case was discharged from a healthcare facility 4–12 weeks before the onset of symptoms.

Data collection: the three options

Data are collected following either the 'minimal', the 'light' or the 'enhanced' CDI surveillance option. As shown in the table below, the minimal surveillance option requires collecting information with only Form H, the light surveillance option requires collecting information with Form H and Form C, and the enhanced surveillance option requires collecting information with Forms H and C as well as Forms E and M.

	Minimal surveillance	Light surveillance	Enhanced surveillance	Form
Collected information	<ul style="list-style-type: none"> • Minimum CDI surveillance for each hospital (aggregated numerator data) • Hospital data for each hospital (aggregated denominator data) 	<ul style="list-style-type: none"> • Minimum CDI surveillance for each hospital (aggregated numerator data) • Hospital data for each hospital (aggregated denominator data) 	<ul style="list-style-type: none"> • Minimum CDI surveillance for each hospital (aggregated numerator data) • Hospital data for each hospital (aggregated denominator data) 	<ul style="list-style-type: none"> • Form H (aggregated numerator and denominator data)
		<ul style="list-style-type: none"> • Information on each CDI case (case-based numerator data) 	<ul style="list-style-type: none"> • Information on each CDI case (case-based numerator data) 	<ul style="list-style-type: none"> • Form C (case-based numerator data)
			<ul style="list-style-type: none"> • Additional information on each CDI case (enhanced numerator data) 	<ul style="list-style-type: none"> • Form E (additional case-based numerator data; one form for each CDI case and <i>C. difficile</i> isolate)
			<ul style="list-style-type: none"> • Microbiological data (for the first 10 consecutively detected cases in each participating healthcare facility: characterisation, susceptibility testing and typing of each <i>C. difficile</i> isolate) 	<ul style="list-style-type: none"> • Form M (one form for each <i>C. difficile</i> isolate)
Surveillance period	<p>Recommended: continuous surveillance for 12 months, starting on the first* day of the month.</p>			
	<p>The recommended minimum surveillance period is three consecutive months, preferably from 1 October to 31 December, or from 1 January to 31 March.</p>			
	<p>Note that on average, a 300-bed European hospital (with 100% bed occupancy) can expect seven CDI cases every three months, or 28 cases per year, for an incidence of three CDI cases per 10 000 patient-days.</p> <p><i>*The pilot study demonstrated that completion of Form H is made much easier by starting surveillance on the first day of a month.</i></p>			

Who collects the data?

The composition of the team responsible for data collection may vary from one hospital to another. It is recommended that hospital infection control personnel as well as the team in charge of the patients are both involved. It is likely that most hospitals using the enhanced surveillance module will acquire microbiological data (Form M) from clinical microbiology laboratory personnel.

Form H: Hospital-based data

This form is used to collect denominator data in the minimal, light and enhanced surveillance options. The minimum requirement for CDI surveillance is completion of Form H alone.

Hospital-based aggregated denominator data are collected for all eligible patients within a participating hospital. One Form H should be filled out for each surveillance period. The recommended minimum surveillance period is three consecutive months, from 1 October to 31 December, or from 1 January to 31 March.

In addition to the denominator data, the following aggregated data are collected for each surveillance period at the hospital level:

- Basic hospital characteristics: hospital type and size, necessary for stratification of incidence rates
- Aggregated numerator data: together with the denominator data, these data allow the calculation of the incidence of healthcare-associated (and total) CDI in participating hospitals, and therefore correspond to the minimal data set for CDI surveillance. The number of cases reported on this form should correspond to the number of completed case files in the light surveillance option.
- Frequency of testing for CDI and diagnostic tests in use: process indicator of surveillance sensitivity.

If a hospital has several facilities located on different sites, data should be only merged for those sites which are related in terms of infection control.

Definitions

Hospital code (required): hospital identifier/code is assigned by the national/regional CDI surveillance coordinator. Hospital codes should be unique within each surveillance network, and kept constant between the ECDC Antimicrobial Resistance and Healthcare-Associated Infection (ARHAI) surveillance protocols and from one year to the next.

Hospital type (required)

Primary	Often referred to as a 'district hospital' or 'first-level referral' hospital.
	Few specialities (mainly internal medicine, obstetrics-gynaecology, paediatrics, general surgery or only general practice).
	Limited laboratory services are available for general, but not for specialised pathological analysis.
	Often corresponds to a general hospital without teaching function.
Secondary	Often referred to as a 'provincial hospital'.
	Hospital is highly differentiated by function with five to ten clinical specialities, such as haematology, oncology, nephrology, ICU.
	Takes some referrals from other (primary) hospitals.
	Often corresponds to a general hospital with teaching function.
Tertiary	Often referred to as a 'central', 'regional' or 'tertiary-level' hospital.
	Highly specialised staff and technical equipment (ICU, haematology, transplantation, cardio-thoracic surgery, neurosurgery); specialised imaging units.
	Clinical services are highly differentiated by function.
	Provides regional services and regularly takes referrals from other (primary and secondary) hospitals.
	Often a university hospital or associated with a university.
Specialised	Single clinical specialty, possibly with sub-specialties.
	Highly specialised staff and technical equipment.
	Specify (e.g. paediatric hospital, infectious diseases hospital), free text.

Surveillance period (required for each surveillance period): start and end date for the CDI surveillance period.

Number of beds (required): number of hospital beds for the current surveillance period. All wards should be included for the surveillance of CDI, exclusion of wards is not allowed. If despite this recommendation certain wards were excluded, it is crucial that the aggregated denominator data are provided for the included wards only.

Number of discharges/admissions (required): number of hospital discharges in the current surveillance period, use number of admissions if discharges are not available.

Number of patient-days (required): number of hospital patient-days in the current surveillance period.

Number of HA CDI cases (required): number of healthcare-associated CDI cases within the surveillance period (i.e. with onset on day three or later, following admission to a healthcare facility on day one, OR in the community within four weeks of discharge from any healthcare facility). Exclude recurrent cases.

Number of CA CDI cases and CDI cases of unknown origin (required): number of community-associated CDI cases and cases of unknown origin within the surveillance period i.e. onset outside of healthcare facilities, AND without discharge from a healthcare facility within the previous 12 weeks, OR onset on the day of admission to a healthcare facility or on the following day AND not resident in a healthcare facility within the previous 12 weeks, OR a CDI case discharged from a healthcare facility 4–12 weeks before the onset. Exclude recurrent cases.

Number of recurrent CDI cases: number of CDI episodes with onset within two and eight weeks of a previous episode (including both healthcare-associated and community-associated recurrent cases).

Number of stool specimens tested: number of stool specimens tested for CDI in the surveillance period. Each specimen should only be counted once, even if more than one test was performed on that specimen.

Number of stool specimens that tested positive for CDI: number of stools tested for CDI with a positive test result in the surveillance period. Each specimen should only be counted once.

Algorithm used for CDI diagnosis: laboratory test(s) applied on faeces samples to recognise the presence of toxin-producing *C. difficile*, either as a solitary test or as a combination of screening and confirmatory tests. If none of the algorithms match your algorithm, indicate the algorithm which matches most closely. If multiple algorithms are applied (i.e. depending on work hours or patient categories), please indicate the most frequently applied algorithm(s), that is/are used for more than 80% of the samples tested for *C. difficile*.

- NAAT: Nucleic Acid Amplification Test
- EIA: Enzyme Immunoassay
- Toxicogenic culture: Culture method for the detection of toxin-producing *C. difficile*; includes testing cultured isolates by a toxin test.



European surveillance of *Clostridium difficile* infections

Form H: Hospital-based data (All types of surveillance)

Hospital code: _____

Hospital type:

- ☐ Primary ☐ Secondary
☐ Tertiary ☐ Specialised hospital; (please specify: _____)

Surveillance period: From ____ / ____ / 20____ (dd/mm/yyyy) to: ____ / ____ / 20____ (dd/mm/yyyy)

For the above surveillance period, specify:

Attribute:	Number
No. of beds	
No. of discharges/admissions	
No. of patient-days	
No. of HA ^{1,3} CDI cases	
No. of CA ^{2,3} CDI cases or CDI cases of unknown origin	
No. of recurrent CDI cases	
No. of stool specimens tested for CDI	
No. of stool specimens that tested positive for CDI	

¹HA: healthcare-associated; ²CA: community-associated; ³recurrent cases excluded

Exclusion of wards/units:

☐ No (recommended) ☐ Yes (not recommended)

If some wards/units were excluded, specify which wards/units were excluded:

Important: All wards/units should be included for the surveillance of CDI. If despite this recommendation certain wards/units were excluded, it is crucial that the aggregated denominator data are provided for the included wards/units only.

Algorithm used for CDI diagnosis:

The diagnostic algorithms below are categorised in decreasing order of expected test validity (maximised sensitivity and specificity). If none of the algorithms below is adequate, indicate the test algorithm which is the closest to the one that you apply. If you apply multiple algorithms, please indicate the most frequently applied algorithm(s), that is/are used for >80% of the samples tested for *C. difficile*. GDH= glutamate dehydrogenase.

Category 1:

- ☐ Screening test with NAAT, confirmation with EIA toxin detection
☐ Screening test with both GDH and toxin detection, confirmation with NAAT
☐ Screening test with both GDH and toxin detection, confirmation with toxigenic culture

Category 2:

- ☐ Screening test with GDH, confirmation with NAAT
☐ Screening test with GDH, confirmation with toxigenic culture
☐ NAAT alone

Category 3:

- ☐ Screening with toxin detection, confirmation with NAAT or toxigenic culture
☐ Toxigenic culture alone
☐ EIA for toxins alone
☐ Stool cytotoxicity assay alone
☐ Multiple methods for the same stool specimen
☐ Other, please specify:

Form C: Case-based data

This form is used to collect numerator data in the light and enhanced surveillance options. Case-based numerator data are collected for all hospitalised patients that meet the CDI case definition and inclusion criteria (see above), including both those with symptoms at admission and those who developed symptoms after admission.

Definitions

Hospital code (required): hospital identifier/code is assigned by the national/regional CDI surveillance coordinator. Hospital codes should be unique within each surveillance network and kept constant between ECDC ARHAI surveillance protocols and from one year to the next.

Surveillance period (required) start and end date for the surveillance in the entire hospital. This will be linked with the denominator data.

Patient counter (required) provide an anonymised patient number. In enhanced surveillance, this number should permit linkage of patient data with microbiological typing/susceptibility data and patient data from enhanced surveillance. Patient identifiers must not be used.

Age in years patient age in years; if missing=unknown (UNK). Provide the patient's age in months if the patient is less than two years old.

Sex. gender of the patient: M (male), F (female).

Date of hospital admission (required) date patient was admitted to the hospital for the current hospitalisation (dd/mm/yyyy).

Recurrent CDI choose yes if the patient had an episode of CDI (return of diarrhoeal stools with a positive laboratory test after the end of treatment) for more than two weeks and less than eight weeks following the onset of a previous episode.

Symptoms of CDI present at admission (required) patient had CDI symptoms when admitted for this episode, Yes/No/Unknown.

Date of onset of CDI symptoms this is mandatory if symptom onset was during current hospitalisation, but not recorded if signs/symptoms were present on admission. Record the date of the first signs or symptoms of the infection (dd/mm/yyyy). If unknown, record the date treatment was started for this infection or the date the first diagnostic sample was taken. If no treatment or sample, please estimate.

Origin: Choose one (for detailed definitions, see Definitions section):

- **Healthcare-associated CDI:** a case with onset of symptoms on day three or later, following admission to a healthcare facility on day one, OR in the community within four weeks of discharge from any healthcare facility. This may apply to the current hospital or a previous stay in another healthcare facility, e.g. in another hospital, a long-term care facility or other healthcare facilities (like outpatient departments etc.).
- **Community-associated CDI:** a case with [onset outside of healthcare facilities, AND without discharge from a healthcare facility within the previous 12 weeks] OR [onset on the day of admission to a healthcare facility or on the following day AND not resident in a healthcare facility within the previous 12 weeks]
- **Unknown association:** a case who was discharged from a healthcare facility 4–12 weeks before symptom onset

Patient outcome: status of the patient at hospital discharge or at end of follow-up in the hospital

- **Discharged alive:** patient was discharged alive; OR patient was still in the hospital and alive at end of follow-up during this hospital stay.
- **Death, CDI definitely contributed to death:** use this category if a causal link between CDI and death can be demonstrated.
- **Death, CDI possibly contributed to death:** use this category if no causal link between CDI and this case's death can be demonstrated, but it is still plausible that CDI was at least a contributory factor.
- **Death, unrelated to CDI:** use this category if the cause of death can be demonstrated not to be related to CDI.
- **Death, relationship to CDI unknown:** use this category if no evidence of contributory factors to the cause of death is available.
- **Unknown:** unknown patient outcome.

Date of discharge/in-hospital death: date the patient was discharged from the hospital; OR date of end of follow-up if the patient was still hospitalised and alive; OR date of death if patient died during the current hospitalisation.

Enhanced data collected for this patient: Yes/No/UNK. If yes, please fill Form E.

Microbiological data collected for this patient: Yes/No/UNK. Indicate whether Form M has been completed.



European surveillance of *Clostridium difficile* infections Form C: Case-based data (Light and enhanced surveillance)

Hospital code: _____

Surveillance period: From ____ / ____ / 20____ (dd/mm/yyyy) to: ____ / ____ / 20____ (dd/mm/yyyy)

Patient counter: _____

Age in years: ____; age if < 2 years old: ____ months.

Sex: ☐ M ☐ F

Date of hospital admission: ____ / ____ / 20____ (dd/mm/yyyy)

Recurrent CDI (positive laboratory tests for CDI in diarrhoeal stools after the end of treatment for CDI occurring > 2 weeks and < 8 weeks following the onset of a previous episode):

☐ Yes ☐ No ☐ Unknown

Symptoms of CDI present at admission:

☐ Yes ☐ No ☐ Unknown

Date of onset of CDI symptoms: ____ / ____ / 20____ (dd/mm/yyyy)

CDI origin (tick one):

- ☐ **Healthcare-associated** (symptom onset on day three or later following admission to a healthcare facility on day one, OR in the community within 4 weeks following discharge from any healthcare facility)
Specify: ☐ Current hospital ☐ Other hospital
 ☐ Long-term care facility ☐ Other healthcare-associated origin
- ☐ **Community-associated** (symptom onset [outside of healthcare facilities, AND without discharge from a healthcare facility within the previous 12 weeks], OR [on the day of admission to a healthcare facility or on the following day AND no residence in a healthcare facility within the previous 12 weeks])
- ☐ **Unknown association** (including cases discharged from a healthcare facility 4–12 weeks before symptom onset)

Patient outcome:

- ☐ Discharged alive
☐ Death, CDI definitely contributed to death
☐ Death, CDI possibly contributed to death
☐ Death, no relation to CDI
☐ Death, relationship to CDI unknown
☐ Unknown

Date of hospital discharge/in-hospital death (dd/mm/yyyy): ____ / ____ / ____

Enhanced data (Form E) collected for this patient: ☐ Yes ☐ No ☐ Unknown

Microbiological data (Form M) collected for this patient: ☐ Yes ☐ No ☐ Unknown

Form E: Additional case-based data

This form is used in the enhanced surveillance option only. Enhanced case-based data should only be collected together with microbiological data, i.e. for a maximum of 10 consecutive patients with CDI.

Definitions

Hospital code (required): hospital identifier/code assigned by national/regional CDI surveillance coordinator. Hospital codes should be unique within each surveillance network and kept constant between ECDC ARHAI surveillance protocols and from one year to another.

Patient counter (required): provide an anonymised patient number that will permit linkage of patient data and microbiological typing/susceptibility data, and between patient data from light and enhanced surveillance. Patient identifiers must not be used.

Ward/unit ID: abbreviated name of hospital ward; it should be used consistently and should remain the same in different surveillance periods/years.

Ward/unit specialty (see code list): main ward specialty; see specialty code list below.

Previous healthcare admission: previous admission in a healthcare facility in the last three months relative to the onset of CDI: Yes/No/Unknown, if yes: Admission in a hospital or another healthcare facility (long-term care, outpatient department, etc.). Collect from electronic records and/or patient notes, and/or by asking the patient.

Physical status: classification of the severity of underlying medical conditions using the McCabe Score [4]. Answer categories: Non-fatal disease (expected survival at least five years); ultimately fatal disease (expected survival between one and five years); rapidly fatal disease (expected death within one year); unknown. Some examples of the McCabe score categories for different diseases are given below. These are not meant to be exhaustive but rather to serve as a guidance tool as part of this protocol.

McCabe score categories	Examples
Rapidly fatal (< one year)	<ul style="list-style-type: none"> End-stage haematological malignancies (unsuitable for transplant, or relapsed), heart failure (EF < 25%) and end-stage liver disease (unsuitable for transplant with recalcitrant ascites, encephalopathy or varices) Multiple organ failure on intensive care unit – APACHE II score > 30, SAPS II score > 70 Pulmonary disease with cor pulmonale
Ultimately fatal: (one year to four years)	<ul style="list-style-type: none"> Chronic leukaemia's, myelomas, lymphomas, metastatic carcinoma, end-stage kidney disease (without transplant) Motor neuron disease, multiple sclerosis non-responsive to treatment Alzheimer's/dementia Diabetes requiring amputation or post amputation
Non-fatal (> five years)	<ul style="list-style-type: none"> Diabetes Carcinoma/haematological malignancy with > 80% five-year survival Inflammatory disorders Chronic GI, GU conditions Obstetrics Infections (including HIV, HCV, HBV – unless in above categories) All other diseases

EF: Ejection fraction, GI: Gastrointestinal, GU: Genitourinary, HCV: Hepatitis C virus, HBV: Hepatitis B virus

Complicated course of CDI: Yes / No / Unknown. CDI leading to any of the following:

- admission to an intensive care unit for treatment of CDI or its complications (e.g. for shock requiring vasopressor therapy);
- surgery (colectomy) for toxic megacolon, perforation or refractory colitis;
- death within 30 days after diagnosis if CDI is either a primary or contributing cause.



European surveillance of *Clostridium difficile* infections

Form E: Additional case-based data (Enhanced surveillance)

(to be combined with form M and form C)

Hospital code: _____

Patient counter: _____

Ward/unit ID: _____

Ward/unit specialty (see code list): _____

Previous healthcare admission in the last 3 months:

☐ Yes ☐ No ☐ Unknown

If yes:

☐ Hospital ☐ Long-term care facility ☐ Other

Physical status (McCabe Score):

☐ Non-fatal underlying disease (survival at least 5 years)

☐ Ultimately fatal underlying disease (survival 1–4 years)

☐ Rapidly fatal underlying disease (survival <1 year)

☐ Unknown

Complicated course of CDI: (i.e. CDI resulted in e.g. ICU admission, toxic megacolon, surgery or death)

☐ Yes ☐ No ☐ Unknown

Form M: Isolate shipment data sheet

This form is only used in the enhanced surveillance option.

Stool samples from a maximum of 10 consecutive patients with primary or recurrent CDI that tested positive for CDI should be stored at -20°C, and cultured for the presence of toxin-producing *C. difficile* using the standard operating procedure for the culture and identification of *C. difficile* (available on request from ECDC), or national or local protocols. Culture methods should be carried out under containment level 2 conditions using the principle of 'good laboratory practice', or containment level 3 if Hazard Group 3 organisms are suspected to be in the specimen.

C. difficile isolates should be sent for typing and characterisation to a laboratory designated at the national level by the national coordinator, accompanied by a partially filled form M. If typing and characterisation is not available at the national level, support from another laboratory (e.g. as defined by ECDC) should be sought.

Definitions

Network-Id: unique identifier for each surveillance network, selected and generated by Member State, e.g. EN, NI, SC, WA for UK or different CClin networks in France; this field is combined with the hospital identifier to create a unique hospital code since different networks within one country may use the same hospital code. Network ID can be omitted if the hospital identifiers are unique within the reporting country.

Hospital code (required): hospital identifier/code assigned by national/regional CDI surveillance coordinator. Hospital codes should be unique within each surveillance network and kept constant between ECDC ARHAI surveillance protocols and from one year to another.

Laboratory code (required): local laboratory identifier/code assigned by national/regional CDI surveillance coordinating centre. For the primary lab responsible for microbiological confirmation of CDI (not the code of the national/reference laboratory). It is recommended to use the same laboratory codes as in EARS-Net.

Surveillance period (required): start and end date for the CDI surveillance period in the entire hospital.

Patient counter (required): provide an anonymised patient number that will permit linkage of patient data and microbiological typing/susceptibility data, and between patient data from light and enhanced surveillance. Patient identifiers must not be used.

Age in years: patient age in years; if missing=UNK. Provide the patient's age in months if the patient is less than two years old.

PCR ribotype of *C. difficile* isolate: *C. difficile* PCR ribotype as determined by conventional gel-electrophoresis or capillary-PCR ribotyping.

Production of toxins A and/or B: production of toxins as determined by PCR of TcdA and TcdB or by EIA;

Antimicrobial susceptibility testing: MIC (minimum inhibitory concentration) and test used for the determination of the MIC.



European surveillance of *Clostridium difficile* infections

Form M: Isolate shipment data sheet (Enhanced surveillance)

(one form for each isolate)

Network-Id: _____

Hospital code: _____

Laboratory code: _____

Surveillance period: From ____ / ____ / 20____ (dd/mm/yyyy) to: ____ / ____ / 20____ (dd/mm/yyyy)

Patient counter: _____

Age in years: ____ (if <2 years old, age in months: ____).

Microbiological results:

Typing: performed by the national reference laboratory:

☐ Yes ☐ No

PCR ribotype of *C. difficile* isolate: _____

Production of toxins A and/or B

☐ Positive ☐ Negative

Presence of binary toxin genes

☐ Positive ☐ Negative

Antimicrobial susceptibility testing: performed by the national reference laboratory:

☐ Yes ☐ No

Metronidazole MIC: _____ mg/l by (method): _____

Vancomycin MIC: _____ mg/l by (method): _____

Moxifloxacin MIC: _____ mg/l by (method): _____

References

1. Kuijper EJ, Coignard B, Tüll P, et al. Emergence of *Clostridium difficile*-associated disease in North America and Europe. Clin Microbiol Infect 2006; 12 (Suppl 6):2-18.
2. Bauer MP, Notermans DW, van Benthem BH, et al. *Clostridium difficile* infection in Europe: a hospital-based survey. Lancet 2011; 377:63-73.
3. Barbut F, Mastrantonio P, Delmée M, et al. Prospective study of *Clostridium difficile* infections in Europe with phenotypic and genotypic characterisation of the isolates. Clin Microbiol Infect 2007; 13:1048-1057.
4. McCabe RW, Jackson GG. Gram-negative bacteremia I. Etiology and ecology. Arch Intern Med 1962; 110: 847-855.

Annex - Ward/unit specialty code list

Specialty codes used for ward/unit specialty (Form E) and hospital specialisation (Form H).

Category	Code	Specialty
Surgical specialties	SURGEN	General surgery
	SURDIG	Digestive tract surgery
	SURORTR	Orthopaedics and surgical traumatology
	SURORTO	Orthopaedics
	SURTR	Traumatology
	SURCV	Cardio surgery and vascular surgery
	SURCARD	Cardio surgery
	SURVASC	Vascular surgery
	SURTHO	Thoracic surgery
	SURNEU	Neurosurgery
	SURPED	Paediatric general surgery
	SURTRANS	Transplantation surgery
	SURONCO	Surgery for cancer
	SURENT	Ear, nose, throat or otorhinolaryngology
	SUROPH	Ophthalmology
	SURMAXFAC	Maxillo-facial surgery
	SURSTODEN	Stomatology/Dentistry
	SURBURN	Burns care
	SURURO	Urology
	SURPLAS	Plastic and reconstructive surgery
	SUROTH	Other surgery
Medical specialties	MEDGEN	General medicine
	MEDGAST	Gastro-enterology
	MEDHEP	Hepatology
	MEDENDO	Endocrinology
	MEDONCO	Oncology
	MEDHEMA	Haematology
	MEDBMT	Bone marrow transplantation (BMT)
	MEDHEMBMT	Haematology/BMT
	MEDCARD	Cardiology
	MEDDERM	Dermatology
	MEDNEPH	Nephrology
	MEDNEU	Neurology
	MEDPNEU	Pneumology
	MEDRHEU	Rheumatology
	MEDID	Infectious diseases
	MEDTR	Medical traumatology
	MEDOTH	Other medical
Paediatrics	PEDNEO	Neonatology
	PEDGEN	Paediatrics general, not specialised
	PEDBAB	Healthy babies on paediatric ward
Intensive care medicine	ICUMED	Medical ICU
	ICUSUR	Surgical ICU
	ICUPED	Paediatric ICU
	ICUNEO	Neonatal ICU
	ICUMIX	Mixed (polyvalent) ICU, general intensive or critical care
	ICUSPEC	Specialised ICU
	ICUOTH	Other ICU
Gynaecology/Obstetrics	GOOBS	Obstetrics/maternity
	GOGYN	Gynaecology
	GOBAB	Healthy babies on maternity ward
Geriatrics	GER	Geriatrics, care for the elderly

Category	Code	Specialty
Psychiatrics	PSY	Psychiatrics
Rehabilitation	RHB	Rehabilitation
Long-term care	LTC	Long-term care
Other	OTH	Other specialty, not listed
Mixed	MIX	Combination of specialties