
Annex 2

WHO Model List of Essential Medicines for Children (April 2015)

Explanatory notes

This Model List is intended for use for children up to 12 years of age.

The **core list** presents a list of minimum medicine needs for a basic health-care system, listing the most efficacious, safe and cost-effective medicines for priority conditions. Priority conditions are selected on the basis of current and estimated future public health relevance, and potential for safe and cost-effective treatment.

The **complementary list** presents essential medicines for priority diseases, for which specialized diagnostic or monitoring facilities, and/or specialist medical care, and/or specialist training are needed. In case of doubt medicines may also be listed as complementary on the basis of consistent higher costs or less attractive cost-effectiveness in a variety of settings.

The **square box symbol** (□) is primarily intended to indicate similar clinical performance within a pharmacological class. The listed medicine should be the example of the class for which there is the best evidence for effectiveness and safety. In some cases, this may be the first medicine that is licensed for marketing; in other instances, subsequently licensed compounds may be safer or more effective. Where there is no difference in terms of efficacy and safety data, the listed medicine should be the one that is generally available at the lowest price, based on international drug price information sources.

Therapeutic equivalence is indicated only on the basis of reviews of efficacy and safety and when consistent with WHO clinical guidelines. National lists should not use a similar symbol and should be specific in their final selection, which would depend on local availability and price.

The format and numbering of the 18th WHO Model List of Essential Medicines have been retained but, as indicated in the text, some sections have been deleted because they contain medicines that are not relevant for children.

a indicates that there is an age or weight restriction on use of the medicines; the details for each medicine are in Table 1.1 of Annex 1.

The presence of an entry on the Essential Medicines List carries no assurance as to pharmaceutical quality. It is the responsibility of the relevant national or regional drug regulatory authority to ensure that each product is of appropriate pharmaceutical quality (including stability) and that when relevant, different products are interchangeable.

For recommendations and advice concerning all aspects of the quality assurance of medicines see the WHO Medicines web site http://www.who.int/medicines/areas/quality_assurance.

Medicines and dosage forms are listed in alphabetical order within each section and there is no implication of preference for one form over another. Standard treatment guidelines should be consulted for information on appropriate dosage forms.

The main terms used for dosage forms in the Essential Medicines List can be found in Table 1.2 of Annex 1.




Definitions of many of these terms and pharmaceutical quality requirements applicable to the different categories are published in the current edition of *The International Pharmacopoeia* <http://www.who.int/medicines/publications/pharmacopoeia>.

1. ANAESTHETICS	
1.1 General anaesthetics and oxygen	
1.1.1 Inhalational medicines	
halothane	Inhalation.
isoflurane	Inhalation.
nitrous oxide	Inhalation.
oxygen	Inhalation (medicinal gas).
1.1.2 Injectable medicines	
ketamine	Injection: 50 mg (as hydrochloride)/mL in 10-mL vial.
propofol *	Injection: 10 mg/mL; 20 mg/mL. * Thiopental may be used as an alternative depending on local availability and cost.
1.2 Local anaesthetics	
<input type="checkbox"/> bupivacaine	Injection: 0.25%; 0.5% (hydrochloride) in vial. Injection for spinal anaesthesia: 0.5% (hydrochloride) in 4-mL ampoule to be mixed with 7.5% glucose solution.
<input type="checkbox"/> lidocaine	Injection: 1%; 2% (hydrochloride) in vial. Injection for spinal anaesthesia: 5% (hydrochloride) in 2-mL ampoule to be mixed with 7.5% glucose solution. Topical forms: 2% to 4% (hydrochloride).
lidocaine + epinephrine (adrenaline)	Dental cartridge: 2% (hydrochloride) + epinephrine 1:80 000. Injection: 1%; 2% (hydrochloride or sulfate) + epinephrine 1:200 000 in vial.
1.3 Preoperative medication and sedation for short-term procedures	
atropine	Injection: 1 mg (sulfate) in 1-mL ampoule.
<input type="checkbox"/> midazolam	Injection: 1 mg/mL. Oral liquid: 2 mg/mL. Tablet: 7.5 mg; 15 mg.
morphine	Injection: 10 mg (sulfate or hydrochloride) in 1-mL ampoule.
2. MEDICINES FOR PAIN AND PALLIATIVE CARE	
2.1 Non-opioids and non-steroidal anti-inflammatory medicines (NSAIMs)	
ibuprofen <input type="checkbox"/>	Oral liquid: 200 mg/5 mL. Tablet: 200 mg; 400 mg; 600 mg. <input type="checkbox"/> Not in children less than 3 months.
paracetamol*	Oral liquid: 125 mg/5 mL.

	<p>Suppository: 100 mg.</p> <p>Tablet: 100 mg to 500 mg.</p> <p>* Not recommended for anti-inflammatory use due to lack of proven benefit to that effect.</p>
2.2 Opioid analgesics	
□ morphine*	<p>Granules (slow release; to mix with water): 20 mg to 200 mg (morphine sulfate).</p> <p>Injection: 10 mg (morphine hydrochloride or morphine sulfate) in 1-mL ampoule.</p> <p>Oral liquid: 10 mg (morphine hydrochloride or morphine sulfate)/5 mL.</p> <p>Tablet (slow release): 10 mg – 200mg (morphine hydrochloride or morphine sulfate).</p> <p>Tablet (immediate release): 10 mg (morphine sulfate).</p> <p>*Alternatives limited to hydromorphone and oxycodone.</p>
2.3 Medicines for other symptoms common in palliative care	
amitriptyline	Tablet: 10 mg; 25 mg.
cyclizine	Injection: 50 mg/mL. Tablet: 50 mg.
dexamethasone	Injection: 4 mg/mL in 1-mL ampoule (as disodium phosphate salt). Oral liquid: 2 mg/5 mL. Tablet: 2 mg.
diazepam	Injection: 5 mg/mL. Oral liquid: 2 mg/5 mL. Rectal solution: 2.5 mg; 5 mg; 10 mg. Tablet: 5 mg; 10 mg.
docusate sodium	Capsule: 100 mg. Oral liquid: 50 mg/5 mL.
fluoxetine a	Solid oral dosage form: 20 mg (as hydrochloride). a >8 years.
hyoscine hydrobromide	Injection: 400 micrograms/mL; 600 micrograms/mL. Transdermal patches: 1 mg/72 hours.
lactulose	Oral liquid: 3.1–3.7 g/5 mL.
midazolam	Injection: 1 mg/mL; 5 mg/mL. Oral liquid: 2mg/mL. Solid oral dosage form: 7.5 mg; 15 mg.
ondansetron a	Injection: 2 mg base/mL in 2-mL ampoule (as

	hydrochloride). Oral liquid: 4 mg base/5 mL. Solid oral dosage form: Eq 4 mg base; Eq 8 mg base. <input type="checkbox"/> > 1 month.
senna	Oral liquid: 7.5 mg/5 mL.
3. ANTIALLERGICS AND MEDICINES USED IN ANAPHYLAXIS	
dexamethasone	Injection: 4 mg/mL in 1-mL ampoule (as disodium phosphate salt).
epinephrine (adrenaline)	Injection: 1 mg (as hydrochloride or hydrogen tartrate) in 1-mL ampoule.
hydrocortisone	Powder for injection: 100 mg (as sodium succinate) in vial.
<input type="checkbox"/> loratadine *	Oral liquid: 1 mg/mL. Tablet: 10 mg. *There may be a role for sedating antihistamines for limited indications.
<input type="checkbox"/> prednisolone	Oral liquid: 5 mg/mL. Tablet: 5 mg; 25 mg.
4. ANTIDOTES AND OTHER SUBSTANCES USED IN POISONINGS	
4.1 Non-specific	
charcoal, activated	Powder.
4.2 Specific	
acetylcysteine	Injection: 200 mg/mL in 10-mL ampoule. Oral liquid: 10%; 20%.
atropine	Injection: 1 mg (sulfate) in 1-mL ampoule.
calcium gluconate	Injection: 100 mg/mL in 10-mL ampoule.
naloxone	Injection: 400 micrograms (hydrochloride) in 1-mL ampoule.
<i>Complementary List</i>	
<i>deferoxamine</i>	Powder for injection: 500 mg (mesilate) in vial.
<i>dimercaprol</i>	Injection in oil: 50 mg/mL in 2-mL ampoule.
<i>fomepizole</i>	Injection: 5 mg/mL (sulfate) in 20-mL ampoule or 1 g/mL (base) in 1.5-mL ampoule.
<i>sodium calcium edetate</i>	Injection: 200 mg/mL in 5-mL ampoule.
<i>succimer</i>	Solid oral dosage form: 100 mg.
5. ANTICONSULSANTS/ANTIPILEPTICS	
carbamazepine	Oral liquid: 100 mg/5 mL.

	<p>Tablet (chewable): 100 mg; 200 mg.</p> <p>Tablet (scored): 100 mg; 200 mg.</p>
diazepam	Gel or rectal solution: 5 mg/mL in 0.5 mL; 2-mL; 4-mL tubes.
<input type="checkbox"/> lorazepam	Parenteral formulation: 2 mg/mL in 1-mL ampoule; 4 mg/mL in 1-mL ampoule.
midazolam	<p>Solution for oromucosal administration: 5 mg/mL; 10 mg/mL</p> <p>Ampoule*: 1 mg/ mL; 10 mg/mL</p> <p>*for buccal administration when solution for oromucosal administration is not available</p>
phenobarbital	<p>Injection: 200 mg/mL (sodium).</p> <p>Oral liquid: 15 mg/5 mL.</p> <p>Tablet: 15 mg to 100 mg.</p>
phenytoin	<p>Injection: 50 mg/mL in 5-mL vial (sodium salt).</p> <p>Oral liquid: 25 mg to 30 mg/5 mL.*</p> <p>Solid oral dosage form: 25 mg; 50 mg; 100 mg (sodium salt).</p> <p>Tablet (chewable): 50 mg.</p> <p>* The presence of both 25 mg/5 mL and 30 mg/5 mL strengths on the same market would cause confusion in prescribing and dispensing and should be avoided.</p>
valproic acid (sodium valproate)	<p>Oral liquid: 200 mg/5 mL.</p> <p>Tablet (crushable): 100 mg.</p> <p>Tablet (enteric-coated): 200 mg; 500 mg (sodium valproate).</p>
Complementary List	
<i>ethosuximide</i>	<p>Capsule: 250 mg.</p> <p>Oral liquid: 250 mg/5 mL.</p>
<i>valproic acid (sodium valproate)</i>	Injection: 100 mg/ mL in 4- mL ampoule; 100 mg/ mL in 10-mL ampoule.
6. ANTI-INFECTIVE MEDICINES	
6.1 Anthelmintics	
6.1.1 Intestinal anthelmintics	
albendazole	Tablet (chewable): 400 mg.
levamisole	Tablet: 50 mg; 150 mg (as hydrochloride).
mebendazole	Tablet (chewable): 100 mg; 500 mg.
niclosamide	Tablet (chewable): 500 mg.

praziquantel	Tablet: 150 mg; 600 mg.
pyrantel	Oral liquid: 50 mg (as embonate or pamoate)/mL. Tablet (chewable): 250 mg (as embonate or pamoate).
6.1.2 Antifilarials	
albendazole	Tablet (chewable): 400 mg.
diethylcarbamazine	Tablet: 50 mg; 100 mg (dihydrogen citrate).
ivermectin	Tablet (scored): 3 mg.
6.1.3 Antischistosomal and other antitremaatode medicines	
praziquantel	Tablet: 600 mg.
triclabendazole	Tablet: 250 mg.
<i>Complementary List</i>	
<i>oxamniquine*</i>	<i>Capsule:</i> 250 mg. <i>Oral liquid:</i> 250 mg/5 mL. <i>* Oxamniquine is listed for use when praziquantel treatment fails.</i>
6.2 Antibacterials	
6.2.1 Beta-lactam medicines	
amoxicillin	Powder for oral liquid: 125 mg (as trihydrate)/5 mL; 250 mg (as trihydrate)/5 mL. Solid oral dosage form: 250 mg; 500 mg (as trihydrate).
amoxicillin + clavulanic acid	Oral liquid: 125 mg amoxicillin + 31.25 mg clavulanic acid/5 mL AND 250 mg amoxicillin + 62.5 mg clavulanic acid/5 mL. Tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt).
ampicillin	Powder for injection: 500 mg; 1 g (as sodium salt) in vial.
benzathine benzylpenicillin	Powder for injection: 900 mg benzylpenicillin (= 1.2 million IU) in 5-mL vial; 1.44 g benzylpenicillin (= 2.4 million IU) in 5-mL vial.
benzylpenicillin	Powder for injection: 600 mg (= 1 million IU); 3 g (= 5 million IU) (sodium or potassium salt) in vial.
cefalexin	Powder for reconstitution with water: 125 mg/5 mL; 250 mg/5 mL (anhydrous). Solid oral dosage form: 250 mg (as monohydrate).
<input type="checkbox"/> cefazolin* 	Powder for injection: 1 g (as sodium salt) in vial. * For surgical prophylaxis.  > 1 month.
ceftriaxone* 	Powder for injection: 250 mg; 1 g (as sodium salt) in vial.

	<p>* Do not administer with calcium and avoid in infants with hyperbilirubinaemia.</p> <p>a >41 weeks corrected gestational age.</p>
☐ cloxacillin	<p>Capsule: 500 mg; 1 g (as sodium salt).</p> <p>Powder for injection: 500 mg (as sodium salt) in vial.</p> <p>Powder for oral liquid: 125 mg (as sodium salt)/5 mL.</p>
phenoxymethylpenicillin	<p>Powder for oral liquid: 250 mg (as potassium salt)/5 mL.</p> <p>Tablet: 250 mg (as potassium salt).</p>
procaine benzylpenicillin*	<p>Powder for injection: 1 g (=1 million IU); 3 g (=3 million IU) in vial.</p> <p>* Procaine benzylpenicillin is not recommended as first-line treatment for neonatal sepsis except in settings with high neonatal mortality, when given by trained health workers in cases where hospital care is not achievable.</p>
Complementary List	
cefotaxime*	<p>Powder for injection: 250 mg per vial (as sodium salt).</p> <p>* 3rd generation cephalosporin of choice for use in hospitalized neonates.</p>
ceftazidime	<p>Powder for injection: 250 mg or 1 g (as pentahydrate) in vial.</p>
imipenem* + cilastatin*	<p>Powder for injection: 250 mg (as monohydrate) + 250 mg (as sodium salt); 500 mg (as monohydrate) + 500 mg (as sodium salt) in vial.</p> <p>* Only listed for the treatment of life-threatening hospital-based infection due to suspected or proven multidrug-resistant infection. Meropenem is indicated for the treatment of meningitis and is licensed for use in children over the age of 3 months.</p>
6.2.2 Other antibacterials	
azithromycin*	<p>Capsule: 250 mg; 500 mg (anhydrous).</p> <p>Oral liquid: 200 mg/5 mL.</p> <p>* Listed only for trachoma.</p>
chloramphenicol	<p>Capsule: 250 mg.</p> <p>Oily suspension for injection*: 0.5 g (as sodium succinate)/mL in 2-mL ampoule.</p> <p>* Only for the presumptive treatment of epidemic meningitis in children older than 2 years.</p> <p>Oral liquid: 150 mg (as palmitate)/5 mL.</p> <p>Powder for injection: 1 g (sodium succinate) in vial.</p>

ciprofloxacin	Oral liquid: 250 mg/5 mL (anhydrous). Solution for IV infusion: 2 mg/mL (as hyclate). Tablet: 250 mg (as hydrochloride).
doxycycline ^a	Oral liquid: 25 mg/5 mL; 50 mg/5 mL (anhydrous). Solid oral dosage form: 50 mg; 100 mg (as hyclate). ^a Use in children <8 years only for life-threatening infections when no alternative exists.
erythromycin	Powder for oral liquid: 125 mg/5 mL (as stearate or estolate or ethyl succinate). Solid oral dosage form: 250 mg (as stearate or estolate or ethyl succinate).
<input type="checkbox"/> gentamicin	Injection: 10 mg; 40 mg (as sulfate)/mL in 2-mL vial.
metronidazole	Injection: 500 mg in 100-mL vial. Oral liquid: 200 mg (as benzoate)/5 mL. Tablet: 200 mg to 500 mg.
nitrofurantoin	Oral liquid: 25 mg/5 mL. Tablet: 100 mg.
sulfamethoxazole + trimethoprim	Injection: 80 mg + 16 mg/mL in 5-mL ampoule; 80 mg + 16 mg/mL in 10-mL ampoule. Oral liquid: 200 mg + 40 mg/5 mL. Tablet: 100 mg + 20 mg; 400 mg + 80 mg.
trimethoprim ^a	Oral liquid: 50 mg/5 mL. Tablet: 100 mg; 200 mg. ^a >6 months.
Complementary List	
<i>clindamycin</i>	Capsule: 150 mg (as hydrochloride). Injection: 150 mg (as phosphate)/mL. Oral liquid: 75 mg/5 mL (as palmitate).
<i>vancomycin</i>	Powder for injection: 250 mg (as hydrochloride) in vial.
6.2.3 Antileprosy medicines	
Medicines used in the treatment of leprosy should never be used except in combination. Combination therapy is essential to prevent the emergence of drug resistance. Colour-coded blister packs (MDT blister packs) containing standard two-medicine (paucibacillary leprosy) or three-medicine (multibacillary leprosy) combinations for adult and childhood leprosy should be used. MDT blister packs can be supplied free of charge through WHO.	
clofazimine	Capsule: 50 mg; 100 mg.
dapsone	Tablet: 25 mg; 50 mg; 100 mg.
rifampicin	Solid oral dosage form: 150 mg; 300 mg.

6.2.4 Antituberculosis medicines	
WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products and paediatric dosage forms of assured pharmaceutical quality.	
ethambutol	Oral liquid: 25 mg/mL. Tablet: 100 mg; 400 mg (hydrochloride).
isoniazid	Oral liquid: 50 mg/5 mL. Tablet: 100 mg to 300 mg. Tablet (scored): 50 mg.
pyrazinamide	Oral liquid: 30 mg/mL. Tablet: 400 mg. Tablet (dispersible): 150 mg. Tablet (scored): 150 mg.
rifampicin	Oral liquid: 20 mg/mL. Solid oral dosage form: 150 mg; 300 mg.
rifapentine*	Tablet: 150 mg *For treatment of latent TB infection (LTBI) only
Complementary List	
Reserve second-line drugs for the treatment of multidrug-resistant tuberculosis (MDR-TB) should be used in specialized centres adhering to WHO standards for TB control.	
<i>amikacin</i>	Powder for injection: 100 mg; 500 mg; 1 g (as sulfate) in vial.
<i>capreomycin</i>	Powder for injection: 1 g (as sulfate) in vial.
<i>cycloserine</i>	Solid oral dosage form: 250 mg.
<i>ethionamide*</i>	Tablet: 125 mg; 250 mg. *Protionamide may be used as an alternative.
<i>kanamycin</i>	Powder for injection: 1 g (as sulfate) in vial.
<i>levofloxacin*</i>	Tablet: 250 mg; 500 mg. * Ofloxacin and moxifloxacin may be used as alternatives based on availability and programme considerations.
<i>linezolid</i>	Injection for intravenous administration: 2 mg/ mL in 300 mL bag Powder for oral liquid: 100 mg/5 mL, Tablet: 400 mg; 600 mg
<i>p-aminosalicylic acid</i>	Granules: 4 g in sachet. Tablet: 500 mg.
<i>streptomycin</i>	Powder for injection: 1 g (as sulfate) in vial.
6.3 Antifungal medicines	

amphotericin B	Powder for injection: 50 mg in vial (as sodium deoxycholate or liposomal complex).
<input type="checkbox"/> fluconazole	Capsule: 50 mg. Injection: 2 mg/mL in vial. Oral liquid: 50 mg/5 mL.
flucytosine	Capsule: 250 mg. Infusion: 2.5 g in 250 mL.
griseofulvin	Oral liquid: 125 mg/5 mL. Solid oral dosage form: 125 mg; 250 mg.
nystatin	Lozenge: 100 000 IU. Oral liquid: 50 mg/5 mL; 100 000 IU/mL. Tablet: 100 000 IU; 500 000 IU.
<i>Complementary List</i>	
<i>potassium iodide</i>	<i>Saturated solution.</i>
6.4 Antiviral medicines	
<i>6.4.1 Antiherpes medicines</i>	
aciclovir	Oral liquid: 200 mg/5 mL. Powder for injection: 250 mg (as sodium salt) in vial. Tablet: 200 mg.
<i>6.4.2 Antiretrovirals</i>	
<p>Based on current evidence and experience of use, medicines in the following three classes of antiretrovirals are included as essential medicines for treatment and prevention of HIV (prevention of mother-to-child transmission and post-exposure prophylaxis). WHO emphasizes the importance of using these products in accordance with global and national guidelines. WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products and paediatric dosage forms of assured pharmaceutical quality.</p> <p>Scored tablets can be used in children and therefore can be considered for inclusion in the listing of tablets, provided that adequate quality products are available.</p>	
<i>6.4.2.1 Nucleoside/Nucleotide reverse transcriptase inhibitors</i>	
abacavir (ABC)	Oral liquid: 100 mg (as sulfate)/5 mL.
lamivudine (3TC)	Oral liquid: 50 mg/5 mL. Tablet: 150 mg.
stavudine (d4T)	Capsule: 15 mg; 20 mg; 30 mg. Powder for oral liquid: 5 mg/5 mL.
zidovudine (ZDV or AZT)	Capsule: 100 mg. Oral liquid: 50 mg/5 mL.
<i>6.4.2.2 Non-nucleoside reverse transcriptase inhibitors</i>	

efavirenz (EFV or EFZ) ^a	Capsule: 50 mg; 100 mg; 200 mg. Tablet: 200 mg (scored). ^a >3 years or >10 kg.
nevirapine (NVP) ^a	Oral liquid: 50 mg/5 mL. Tablet: 50 mg (dispersible); 200 mg. ^a > 6 weeks
6.4.2.3 Protease inhibitors	
Selection of protease inhibitor(s) from the Model List will need to be determined by each country after consideration of international and national treatment guidelines and experience. Ritonavir is recommended for use in combination as a pharmacological booster, and not as an antiretroviral in its own right. All other protease inhibitors should be used in boosted forms (e.g. with ritonavir).	
atazanavir ^a	Solid oral dosage form: 100 mg; 150 mg (as sulfate). ^a >25 kg.
darunavir ^a	Tablet: 75 mg; ^a >3 years
lopinavir + ritonavir (LPV/r)	Oral liquid: 400 mg + 100 mg/5 mL. Tablet (heat stable): 100 mg + 25 mg-
ritonavir	Oral liquid: 400 mg/5 mL. Tablet (heat stable): 25 mg; 100 mg.
FIXED-DOSE COMBINATIONS	
abacavir + lamivudine	Tablet (dispersible, scored): 60 mg (as sulfate) + 30 mg
lamivudine + nevirapine + stavudine	Tablet (dispersible): 30 mg + 50 mg + 6 mg.
lamivudine + nevirapine + zidovudine	Tablet: 30 mg + 50 mg + 60 mg.
lamivudine + zidovudine	Tablet: 30 mg + 60 mg.
6.4.3 Other antivirals	
oseltamivir*	Capsule: 30 mg; 45 mg; 75 mg (as phosphate). Oral powder: 12 mg/mL. * potentially severe or complicated illness due to confirmed or suspected influenza virus infection in accordance with WHO treatment guidelines.
ribavirin*	Injection for intravenous administration: 800 mg and 1 g in 10-mL phosphate buffer solution. Solid oral dosage form: 200 mg; 400 mg; 600 mg. * For the treatment of viral haemorrhagic fevers only.
6.4.4 Antihepatitis medicines	
6.4.4.1 Medicines for hepatitis B	
entecavir	Oral liquid: 0.05 mg/ mL

	Tablet: 0.5 mg; 1 mg
6.5 Antiprotozoal medicines	
6.5.1 Antiamoebic and anti giardiasis medicines	
diloxanide ^a	Tablet: 500 mg (furoate). ^a >25 kg.
□ metronidazole	Injection: 500 mg in 100-mL vial. Oral liquid: 200 mg (as benzoate)/5 mL. Tablet: 200 mg to 500 mg.
6.5.2 Antileishmaniasis medicines	
amphotericin B	Powder for injection: 50 mg in vial. As sodium deoxycholate or liposomal complex.
miltefosine	Solid oral dosage form: 10 mg; 50 mg.
paromomycin	Solution for intramuscular injection: 750 mg of paromomycin base (as the sulfate).
sodium stibogluconate or meglumine antimoniate	Injection: 100 mg/mL, 1 vial = 30 mL or 30%, equivalent to approximately 8.1% antimony (pentavalent) in 5-mL ampoule.
6.5.3 Antimalarial medicines	
6.5.3.1 For curative treatment	
Medicines for the treatment of <i>P. falciparum</i> malaria cases should be used in combination. The list currently recommends combinations according to treatment guidelines. WHO recognizes that not all of the fixed dose combinations (FDCs in the WHO treatment guidelines exist, and encourages their development and rigorous testing. WHO also encourages development and testing of rectal dosage formulations.	
amodiaquine*	Tablet: 153 mg or 200 mg (as hydrochloride). * To be used in combination with artesunate 50 mg.
artemether*	Oily injection: 80 mg/mL in 1-mL ampoule. * For use in the management of severe malaria.
artemether + lumefantrine*	Tablet: 20 mg + 120 mg. Tablet (dispersible): 20 mg + 120 mg. * Not recommended in the first trimester of pregnancy or in children below 5 kg.
artesunate*	Injection: ampoules, containing 60 mg anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution. For use in the management of severe malaria. Rectal dosage form: 50 mg; 200 mg capsules (for pre-referral treatment of severe malaria only; patients should be taken to an appropriate health facility for follow-up care). Tablet: 50 mg.

	* To be used in combination with either amodiaquine, mefloquine or sulfadoxine + pyrimethamine.
artesunate + amodiaquine *	Tablet: 25 mg + 67.5 mg; 50 mg + 135 mg; 100 mg + 270 mg. * Other combinations that deliver the target doses required such as 153 mg or 200 mg (as hydrochloride) with 50 mg artesunate can be alternatives.
artesunate + mefloquine	Tablet: 25 mg + 55 mg; 100 mg + 220 mg.
chloroquine*	Oral liquid: 50 mg (as phosphate or sulfate)/5 mL. Tablet: 100 mg; 150 mg (as phosphate or sulfate). * For use only for the treatment of <i>P.vivax</i> infection.
doxycycline*	Capsule: 100 mg (as hydrochloride or hyclate). Tablet (dispersible): 100 mg (as monohydrate). * For use only in combination with quinine.
mefloquine*	Tablet: 250 mg (as hydrochloride). * To be used in combination with artesunate 50 mg.
primaquine*	Tablet: 7.5 mg; 15 mg (as diphosphate). * Only for use to achieve radical cure of <i>P.vivax</i> and <i>P.ovale</i> infections, given for 14 days.
quinine*	Injection: 300 mg quinine hydrochloride/mL in 2-mL ampoule. Tablet: 300 mg (quinine sulfate) or 300 mg (quinine bisulfate). * For use only in the management of severe malaria, and should be used in combination with doxycycline.
sulfadoxine + pyrimethamine*	Tablet: 500 mg + 25 mg. * Only in combination with artesunate 50 mg.
6.5.3.2 For prophylaxis	
chloroquine*	Oral liquid: 50 mg (as phosphate or sulfate)/5 mL. Tablet: 150 mg (as phosphate or sulfate). * For use only for the treatment of <i>P.vivax</i> infection.
doxycycline a	Solid oral dosage form: 100 mg (as hydrochloride or hyclate). a >8 years.
mefloquine a	Tablet: 250 mg (as hydrochloride). a >5 kg or >3 months.
proguanil*	Tablet: 100 mg (as hydrochloride). * For use only in combination with chloroquine.
6.5.4 Antipneumocystosis and antitoxoplasmosis medicines	
pyrimethamine	Tablet: 25 mg.

sulfadiazine	Tablet: 500 mg.
sulfamethoxazole + trimethoprim	Injection: 80 mg + 16 mg/mL in 5-mL ampoule; 80 mg + 16 mg/mL in 10-mL ampoule. Oral liquid: 200 mg + 40 mg/5 mL. Tablet: 100 mg + 20 mg; 400 mg + 80 mg.
6.5.5 Antitrypanosomal medicines	
6.5.5.1 African trypanosomiasis	
Medicines for the treatment of 1st stage African trypanosomiasis.	
pentamidine*	Powder for injection: 200 mg (as isetionate) in vial. * To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
suramin sodium*	Powder for injection: 1 g in vial. * To be used for the treatment of the initial phase of <i>Trypanosoma brucei rhodesiense</i> infection.
Medicines for the treatment of 2nd stage African trypanosomiasis	
eflornithine*	Injection: 200 mg (hydrochloride)/mL in 100-mL bottle. * To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
nifurtimox*	Tablet: 120 mg. * Only to be used in combination with eflornithine, for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
Complementary List	
<i>melarsoprol</i>	Injection: 3.6% solution in 5-mL ampoule (180 mg of active compound).
6.5.5.2 American trypanosomiasis	
benznidazole	Tablet: 12.5 mg; 100 mg. Tablet (scored): 50 mg.
nifurtimox	Tablet: 30 mg; 120 mg; 250 mg.
7. ANTIMIGRAINE MEDICINES	
7.1 For treatment of acute attack	
ibuprofen	Tablet: 200 mg; 400 mg.
paracetamol	Oral liquid: 125 mg/5 mL. Tablet: 300 mg to 500 mg.
7.2 For prophylaxis	
propranolol	Tablet: 20 mg; 40 mg (hydrochloride).
8. ANTINEOPLASTICS AND IMMUNOSUPPRESSIVES	
8.1 Immunosuppressive medicines	

Complementary List	
<i>azathioprine</i>	Powder for injection: 100 mg (as sodium salt) in vial. Tablet (scored): 50 mg.
<i>ciclosporin</i>	Capsule: 25 mg. Concentrate for injection: 50 mg/mL in 1-mL ampoule for organ transplantation.
8.2 Cytotoxic and adjuvant medicines	
Medicines listed below should be used according to protocols for treatment of the diseases.	
Complementary List	
<i>asparaginase</i>	Powder for injection: 10 000 IU in vial. – acute lymphoblastic leukemia.
<i>bleomycin</i>	Powder for injection: 15 mg (as sulfate) in vial. – Hodgkin lymphoma – Testicular germ cell tumours – Ovarian germ cell tumours
<i>calcium folinate</i>	Injection: 3 mg/ mL in 10- mL ampoule. Tablet: 15 mg. – Osteosarcoma – Burkitt lymphoma
<i>carboplatin</i>	Injection: 50 mg/5 mL; 150 mg/15 mL; 450 mg/45 mL; 600 mg/60 mL. – osteosarcoma – retinoblastoma
<i>cisplatin</i>	Injection: 50 mg/50 mL; 100 mg/100 mL. – osteosarcoma – testicular germ cell tumours – ovarian germ cell tumours
<i>cyclophosphamide</i>	Powder for injection: 500 mg in vial. Tablet: 25 mg. – rhabdomyosarcoma – Ewing sarcoma – acute lymphoblastic leukemia – Burkitt lymphoma – Hodgkin lymphoma
<i>cytarabine</i>	Powder for injection: 100 mg in vial. – acute lymphoblastic leukemia – Burkitt lymphoma.
<i>dacarbazine</i>	Powder for injection: 100 mg in vial.

	<ul style="list-style-type: none"> – Hodgkin lymphoma
<i>dactinomycin</i>	<p>Powder for injection: 500 micrograms in vial.</p> <ul style="list-style-type: none"> – rhabdomyosarcoma – Wilms tumour
<i>doxorubicin</i>	<p>Powder for injection: 10 mg; 50 mg (hydrochloride) in vial.</p> <ul style="list-style-type: none"> – osteosarcoma – Ewing sarcoma – acute lymphoblastic leukemia – Wilms tumour – Burkitt lymphoma – Hodgkin lymphoma
<i>etoposide</i>	<p>Capsule: 100 mg.</p> <p>Injection: 20 mg/ mL in 5- mL ampoule.</p> <ul style="list-style-type: none"> – retinoblastoma – Ewing sarcoma – acute lymphoblastic leukemia – Burkitt lymphoma – Hodgkin lymphoma – Testicular germ cell tumours – Ovarian germ cell tumours
<i>filgrastim</i>	<p>Injection: 120 micrograms/0.2 mL; 300 micrograms/0.5 mL; 480 micrograms/0.8 mL in pre-filled syringe 300 micrograms/mL in 1- mL vial, 480 mg/1.6 mL in 1.6- mL vial.</p> <ul style="list-style-type: none"> – primary prophylaxis in patients at high risk for developing febrile neutropenia associated with myelotoxic chemotherapy. – secondary prophylaxis for patients who have experienced neutropenia following prior myelotoxic chemotherapy – to facilitate administration of dose dense chemotherapy regimens
<i>ifosfamide</i>	<p>Powder for injection: 500 mg vial 1-g vial; 2-g vial.</p> <ul style="list-style-type: none"> – osteosarcoma – rhabdomyosarcoma – Ewing sarcoma – Testicular germ cell tumours – Ovarian germ cell tumours
<i>mercaptopurine</i>	<p>Tablet: 50 mg.</p> <ul style="list-style-type: none"> – acute lymphoblastic leukemia
<i>mesna</i>	<p>Injection: 100 mg/ mL in 4- mL and 10- mL ampoules.</p> <p>Tablet: 400 mg; 600 mg.</p>

	<ul style="list-style-type: none"> – osteosarcoma – rhabdomyosarcoma – Ewing sarcoma. – Testicular germ cell tumours – Ovarian germ cell tumours
methotrexate	<p>Powder for injection: 50 mg (as sodium salt) in vial.</p> <p>Tablet: 2.5 mg (as sodium salt).</p> <ul style="list-style-type: none"> – osteosarcoma – acute lymphoblastic leukemia
tioguanine [c]	<p>Solid oral dosage form: 40 mg.</p> <ul style="list-style-type: none"> – acute lymphoblastic leukemia.
vinblastine	<p>Powder for injection: 10 mg (sulfate) in vial.</p> <ul style="list-style-type: none"> – Testicular germ cell tumours – Ovarian germ cell tumours
vincristine	<p>Powder for injection: 1 mg; 5 mg (sulfate) in vial.</p> <ul style="list-style-type: none"> – retinoblastoma – rhabdomyosarcoma – Ewing sarcoma – acute lymphoblastic leukemia – Wilms tumour – Burkitt lymphoma. – Hodgkin lymphoma
8.3 Hormones and antihormones	
Complementary List	
dexamethasone	<p>Oral liquid: 2 mg/5 mL</p> <ul style="list-style-type: none"> – acute lymphoblastic leukemia
hydrocortisone	<p>Powder for injection: 100 mg (as sodium succinate) in vial.</p> <ul style="list-style-type: none"> – acute lymphoblastic leukemia.
methylprednisolone [c]	<p>Injection: 40 mg/ mL (as sodium succinate) in 1- mL single-dose vial and 5- mL multi-dose vials; 80 mg/ mL (as sodium succinate) in 1- mL single-dose vial.</p> <ul style="list-style-type: none"> – acute lymphoblastic leukamia.
□ prednisolone	<p>Oral liquid: 5 mg/ mL [c].</p> <p>Tablet: 5 mg; 25 mg.</p> <ul style="list-style-type: none"> – acute lymphoblastic leukemia – Burkitt lymphoma – Hodgkin lymphoma

9. ANTIPARKINSONISM MEDICINES	
10. MEDICINES AFFECTING THE BLOOD	
10.1 Antianaemia medicines	
ferrous salt	Oral liquid: equivalent to 25 mg iron (as sulfate)/mL. Tablet: equivalent to 60 mg iron.
folic acid	Tablet: 1 mg; 5 mg.
hydroxocobalamin	Injection: 1 mg (as acetate, as hydrochloride or as sulfate) in 1-mL ampoule.
10.2 Medicines affecting coagulation	
phytomenadione	Injection: 1 mg/mL; 10 mg/mL in 5-mL ampoule. Tablet: 10 mg.
<i>Complementary List</i>	
<i>desmopressin</i>	Injection: 4 micrograms/ mL (as acetate) in 1- mL ampoule. Nasal spray: 10 micrograms (as acetate) per dose
<i>heparin sodium</i>	Injection: 1000 IU/mL; 5000 IU/mL in 1-mL ampoule.
<i>protamine sulfate</i>	Injection: 10 mg/mL in 5-mL ampoule.
<input type="checkbox"/> <i>warfarin</i>	Tablet: 0.5 mg; 1 mg; 2 mg; 5 mg (sodium salt).
10.3 Other medicines for haemoglobinopathies	
<i>Complementary list</i>	
<i>deferoxamine*</i>	Powder for injection: 500 mg (mesilate) in vial. * <i>Deferasirox oral form may be an alternative, depending on cost and availability.</i>
<i>hydroxycarbamide</i>	Solid oral dosage form: 200 mg; 500 mg; 1 g.
11. BLOOD PRODUCTS OF HUMAN ORIGIN AND PLASMA SUBSTITUTES	
11.1 Blood and blood components	
In accordance with the World Health Assembly resolution WHA63.12, WHO recognizes that achieving self-sufficiency, unless special circumstances preclude it, in the supply of safe blood components based on voluntary, non-remunerated blood donation, and the security of that supply are important national goals to prevent blood shortages and meet the transfusion requirements of the patient population. All preparations should comply with the WHO requirements.	
fresh–frozen plasma	
platelet	
red blood cells	
whole blood	
11.2 Plasma-derived medicines	
All human plasma-derived medicines should comply with the WHO requirements.	
11.2.1 Human immunoglobulins	
anti-rabies immunoglobulin	Injection: 150 IU/ mL in vial.

anti-tetanus immunoglobulin	Injection: 500 IU in vial.
<i>Complementary List</i>	
normal immunoglobulin	Intramuscular administration: 16% protein solution.* Intravenous administration: 5%; 10% protein solution.** Subcutaneous administration: 15%; 16% protein solution.* * Indicated for primary immune deficiency. **Indicated for primary immune deficiency and Kawasaki disease.
11.2.2 Blood coagulation factors	
<i>Complementary List</i>	
<input type="checkbox"/> coagulation factor VIII	Powder for injection: 500 IU/vial.
<input type="checkbox"/> coagulation factor IX	Powder for injection: 500 IU/vial, 1000 IU/vial.
11.3 Plasma substitutes	
<input type="checkbox"/> dextran 70*	Injectable solution: 6%. * Polygeline, injectable solution, 3.5% is considered as equivalent.
12. CARDIOVASCULAR MEDICINES	
12.1 Antianginal medicines	
12.2 Antiarrhythmic medicines	
12.3 Antihypertensive medicines	
<input type="checkbox"/> enalapril	Tablet: 2.5 mg; 5 mg (as hydrogen maleate).
12.4 Medicines used in heart failure	
digoxin	Injection: 250 micrograms/mL in 2-mL ampoule. Oral liquid: 50 micrograms/mL. Tablet: 62.5 micrograms; 250 micrograms.
furosemide	Injection: 10 mg/mL in 2-mL ampoule. Oral liquid: 20 mg/5 mL. Tablet: 40 mg.
<i>Complementary List</i>	
dopamine	Injection: 40 mg (hydrochloride) in 5-mL vial.
12.5 Antithrombotic medicines	
12.6 Lipid-lowering agents	
13. DERMATOLOGICAL MEDICINES (topical)	
13.1 Antifungal medicines	
<input type="checkbox"/> miconazole	Cream or ointment: 2% (nitrate).

terbinafine	Cream: 1% or Ointment: 1% terbinafine hydrochloride.
13.2 Anti-infective medicines	
mupirocin	Cream (as mupirocin calcium): 2%. Ointment: 2%.
potassium permanganate	Aqueous solution: 1:10 000.
silver sulfadiazine <input type="checkbox"/> a	Cream: 1%. a >2 months.
13.3 Anti-inflammatory and antipruritic medicines	
<input type="checkbox"/> betamethasone <input type="checkbox"/> a	Cream or ointment: 0.1% (as valerate). a Hydrocortisone preferred in neonates.
calamine	Lotion.
hydrocortisone	Cream or ointment: 1% (acetate).
13.4 Medicines affecting skin differentiation and proliferation	
benzoyl peroxide	Cream or lotion: 5%.
coal tar	Solution: 5%.
<input type="checkbox"/> podophyllum resin	Solution: 10% to 25%.
salicylic acid	Solution: 5%.
urea	Cream or ointment: 5%; 10%.
13.5 Scabicides and pediculicides	
<input type="checkbox"/> benzyl benzoate <input type="checkbox"/> a	Lotion: 25%. a >2 years.
permethrin	Cream: 5%. Lotion: 1%.
14. DIAGNOSTIC AGENTS	
14.1 Ophthalmic medicines	
fluorescein	Eye drops: 1% (sodium salt).
<input type="checkbox"/> tropicamide	Eye drops: 0.5%.
14.2 Radiocontrast media	
<i>Complementary List</i>	
<i>barium sulfate</i>	<i>Aqueous suspension.</i>
15. DISINFECTANTS AND ANTISEPTICS	
15.1 Antiseptics	
<input type="checkbox"/> chlorhexidine	Solution: 5% (digluconate). Gel: 4%.
<input type="checkbox"/> ethanol	Solution: 70% (denatured).
<input type="checkbox"/> povidone iodine	Solution: 10% (equivalent to 1% available iodine).

15.2 Disinfectants	
alcohol based hand rub	Solution containing ethanol 80% volume /volume Solution containing isopropyl alcohol 75% volume/volume
<input type="checkbox"/> chlorine base compound	Powder: (0.1% available chlorine) for solution.
<input type="checkbox"/> chloroxylenol	Solution: 4.8%.
glutaral	Solution: 2%.
16. DIURETICS	
furosemide	Injection: 10 mg/mL in 2-mL ampoule. Oral liquid: 20 mg/5 mL. Tablet: 10 mg; 20 mg; 40 mg.
<i>Complementary List</i>	
<input type="checkbox"/> hydrochlorothiazide	Tablet (scored): 25 mg.
mannitol	Injectable solution: 10%; 20%.
spironolactone	Oral liquid: 5 mg/5 mL; 10 mg/5 mL; 25 mg/5 mL. Tablet: 25 mg.
17. GASTROINTESTINAL MEDICINES	
<i>Complementary List</i>	
<input type="checkbox"/> pancreatic enzymes	<i>Age-appropriate formulations and doses including lipase, protease and amylase.</i>
17.1 Antiulcer medicines	
<input type="checkbox"/> omeprazole	Powder for oral liquid: 20-mg; 40-mg sachets. Solid oral dosage form: 10 mg; 20 mg; 40 mg.
<input type="checkbox"/> ranitidine	Injection: 25 mg/mL (as hydrochloride) in 2-mL ampoule. Oral liquid: 75 mg/5 mL (as hydrochloride). Tablet: 150 mg (as hydrochloride).
17.2 Antiemetic medicines	
dexamethasone	Injection: 4 mg/mL in 1-mL ampoule (as disodium phosphate salt). Oral liquid: 0.5 mg/5 mL; 2 mg/5 mL. Solid oral dosage form: 0.5 mg; 0.75 mg; 1.5 mg; 4 mg.
metoclopramide ^a	Injection: 5 mg (hydrochloride)/mL in 2-mL ampoule. Oral liquid: 5 mg/5 mL. Tablet: 10 mg (hydrochloride). ^a Not in neonates.

ondansetron ^a	<p>Injection: 2 mg base/mL in 2-mL ampoule (as hydrochloride).</p> <p>Oral liquid: 4 mg base/5 mL.</p> <p>Solid oral dosage form: Eq 4 mg base; Eq 8 mg base.</p> <p>^a >1 month.</p>																				
17.3 Anti-inflammatory medicines																					
17.4 Laxatives																					
17.5 Medicines used in diarrhoea																					
17.5.1 Oral rehydration																					
oral rehydration salts	<p>Powder for dilution in 200 mL; 500 mL; 1 L.</p> <table> <tr> <td>glucose:</td> <td>75 mEq</td> </tr> <tr> <td>sodium:</td> <td>75 mEq or mmol/L</td> </tr> <tr> <td>chloride:</td> <td>65 mEq or mmol/L</td> </tr> <tr> <td>potassium:</td> <td>20 mEq or mmol/L</td> </tr> <tr> <td>citrate:</td> <td>10 mmol/L</td> </tr> <tr> <td>osmolarity:</td> <td>245 mOsm/L</td> </tr> <tr> <td>glucose:</td> <td>13.5 g/L</td> </tr> <tr> <td>sodium chloride:</td> <td>2.6 g/L</td> </tr> <tr> <td>potassium chloride:</td> <td>1.5 g/L</td> </tr> <tr> <td>trisodium citrate dihydrate*:</td> <td>2.9 g/L</td> </tr> </table> <p>* trisodium citrate dihydrate may be replaced by sodium hydrogen carbonate (sodium bicarbonate) 2.5 g/L. However, as the stability of this latter formulation is very poor under tropical conditions, it is recommended only when manufactured for immediate use.</p>	glucose:	75 mEq	sodium:	75 mEq or mmol/L	chloride:	65 mEq or mmol/L	potassium:	20 mEq or mmol/L	citrate:	10 mmol/L	osmolarity:	245 mOsm/L	glucose:	13.5 g/L	sodium chloride:	2.6 g/L	potassium chloride:	1.5 g/L	trisodium citrate dihydrate*:	2.9 g/L
glucose:	75 mEq																				
sodium:	75 mEq or mmol/L																				
chloride:	65 mEq or mmol/L																				
potassium:	20 mEq or mmol/L																				
citrate:	10 mmol/L																				
osmolarity:	245 mOsm/L																				
glucose:	13.5 g/L																				
sodium chloride:	2.6 g/L																				
potassium chloride:	1.5 g/L																				
trisodium citrate dihydrate*:	2.9 g/L																				
17.5.2 Medicines for diarrhoea																					
zinc sulfate*	<p>Solid oral dosage form: 20 mg. ^[c].</p> <p>* In acute diarrhoea zinc sulfate should be used as an adjunct to oral rehydration salts.</p>																				
18. HORMONES, OTHER ENDOCRINE MEDICINES AND CONTRACEPTIVES																					
18.1 Adrenal hormones and synthetic substitutes																					
fludrocortisone	Tablet: 100 micrograms (acetate).																				
hydrocortisone	Tablet: 5 mg; 10 mg; 20 mg.																				
18.2 Androgens																					
18.3 Contraceptives																					
18.3.1 Oral hormonal contraceptives																					
18.3.2 Injectable hormonal contraceptives																					
18.3.3 Intrauterine devices																					
18.3.4 Barrier methods																					
18.3.5 Implantable contraceptives																					

18.4 Estrogens	
18.5 Insulins and other medicines used for diabetes	
glucagon	Injection: 1 mg/mL.
insulin injection (soluble)	Injection: 100 IU/mL in 10-mL vial.
intermediate-acting insulin	Injection: 100 IU/mL in 10-mL vial (as compound insulin zinc suspension or isophane insulin).
<i>Complementary List</i>	
<i>metformin</i>	Tablet: 500 mg (hydrochloride).
18.6 Ovulation inducers	
18.7 Progestogens	
18.8 Thyroid hormones and antithyroid medicines	
levothyroxine	Tablet: 25 micrograms; 50 micrograms; 100 micrograms (sodium salt).
<i>Complementary List</i>	
<i>Lugol's solution</i>	Oral liquid: about 130 mg total iodine/mL.
<i>potassium iodide</i>	Tablet: 60 mg.
<i>propylthiouracil</i>	Tablet: 50 mg.
19. IMMUNOLOGICALS	
19.1 Diagnostic agents	
All tuberculins should comply with the WHO requirements for tuberculins.	
tuberculin, purified protein derivative (PPD)	Injection.
19.2 Sera and immunoglobulins	
All plasma fractions should comply with the WHO requirements.	
Anti-venom immunoglobulin*	Injection. * Exact type to be defined locally.
diphtheria antitoxin	Injection: 10 000 IU; 20 000 IU in vial.
19.3 Vaccines	
<p>WHO immunization policy recommendations are published in vaccine position papers on the basis of recommendations made by the Strategic Advisory Group of Experts on Immunization (SAGE).</p> <p>WHO vaccine position papers are updated three to four times per year. The list below details the vaccines for which there is a recommendation from SAGE and a corresponding WHO position paper as at [DATE]. The most recent versions of the WHO position papers, reflecting the current evidence related to a specific vaccine and the related recommendations, can be accessed at any time on the WHO website at:</p> <p>http://www.who.int/immunization/documents/positionpapers/en/index.html.</p> <p>Vaccine recommendations may be universal or conditional (e.g., in certain regions, in some high-risk populations or as part of immunization programmes with certain characteristics). Details are available in the relevant position papers, and in the Summary Tables of WHO Routine Immunization</p>	

Recommendations available on the WHO website at:

http://www.who.int/immunization/policy/immunization_tables/en/index.html.

Selection of vaccines from the Model List will need to be determined by each country after consideration of international recommendations, epidemiology and national priorities.

All vaccines should comply with the WHO requirements for biological substances.

WHO noted the need for vaccines used in children to be polyvalent.

<i>Recommendations for all</i>	
BCG vaccine	
diphtheria vaccine	
Haemophilus influenzae type b vaccine	
hepatitis B vaccine	
HPV vaccine	
measles vaccine	
pertussis vaccine	
pneumococcal vaccine	
poliomyelitis vaccine	
rotavirus vaccine	
rubella vaccine	
tetanus vaccine	
<i>Recommendations for certain regions</i>	
Japanese encephalitis vaccine	
yellow fever vaccine	
tick-borne encephalitis vaccine	
<i>Recommendations for some high-risk populations</i>	
cholera vaccine	
hepatitis A vaccine	
meningococcal meningitis vaccine	
rabies vaccine	
typhoid vaccine	
<i>Recommendations for immunization programmes with certain characteristics</i>	
influenza vaccine (seasonal)	

mumps vaccine	
varicella vaccine	
20. MUSCLE RELAXANTS (PERIPHERALLY-ACTING) AND CHOLINESTERASE INHIBITORS	
neostigmine	Injection: 500 micrograms in 1-mL ampoule; 2.5 mg (metilsulfate) in 1-mL ampoule. Tablet: 15 mg (bromide).
suxamethonium	Injection: 50 mg (chloride)/mL in 2-mL ampoule. Powder for injection: (chloride), in vial.
<input type="checkbox"/> vecuronium	Powder for injection: 10 mg (bromide) in vial.
<i>Complementary List</i>	
<i>pyridostigmine</i>	Injection: 1 mg in 1-mL ampoule. Tablet: 60 mg (bromide).
21. OPHTHALMOLOGICAL PREPARATIONS	
21.1 Anti-infective agents	
aciclovir	Ointment: 3% W/W.
azithromycin	Solution (eye drops): 1.5%
<input type="checkbox"/> gentamicin	Solution (eye drops): 0.3% (sulfate).
<input type="checkbox"/> ofloxacin	Solution (eye drops): 0.3%.
<input type="checkbox"/> tetracycline	Eye ointment: 1% (hydrochloride).
21.2 Anti-inflammatory agents	
<input type="checkbox"/> prednisolone	Solution (eye drops): 0.5% (sodium phosphate).
21.3 Local anaesthetics	
<input type="checkbox"/> tetracaine ^a	Solution (eye drops): 0.5% (hydrochloride). ^a Not in preterm neonates.
21.4 Miotics and antiglaucoma medicines	
21.5 Mydriatics	
atropine* ^a	Solution (eye drops): 0.1%; 0.5%; 1% (sulfate). * Or homatropine (hydrobromide) or cyclopentolate (hydrochloride). ^a >3 months.
<i>Complementary List</i>	
<i>epinephrine (adrenaline)</i>	Solution (eye drops): 2% (as hydrochloride).
22. OXYTOGICS AND ANTI-OXYTOGICS	
22.1 Oxytocics	
22.2 Antioxytocics (tocolytics)	
23. PERITONEAL DIALYSIS SOLUTION	

<i>Complementary List</i>	
<i>intraperitoneal dialysis solution (of appropriate composition)</i>	<i>Parenteral solution.</i>
24. MEDICINES FOR MENTAL AND BEHAVIOURAL DISORDERS	
24.1 Medicines used in psychotic disorders	
<i>Complementary List</i>	
<i>chlorpromazine</i>	<i>Injection: 25 mg (hydrochloride)/mL in 2-mL ampoule. Oral liquid: 25 mg (hydrochloride)/5 mL. Tablet: 10 mg; 25 mg; 50 mg; 100 mg (hydrochloride).</i>
<i>haloperidol</i>	<i>Injection: 5 mg in 1-mL ampoule. Oral liquid: 2 mg/mL. Solid oral dosage form: 0.5 mg; 2 mg; 5 mg.</i>
24.2 Medicines used in mood disorders	
24.2.1 Medicines used in depressive disorders	
<i>Complementary List</i>	
<i>fluoxetine ^a</i>	<i>Solid oral dosage form: 20 mg (as hydrochloride). ^a >8 years.</i>
24.2.2 Medicines used in bipolar disorders	
24.3 Medicines for anxiety disorders	
24.4 Medicines used for obsessive compulsive disorders	
24.5 Medicines for disorders due to psychoactive substance use	
25. MEDICINES ACTING ON THE RESPIRATORY TRACT	
25.1 Antiasthmatic medicines	
<input type="checkbox"/> budesonide	Inhalation (aerosol): 100 micrograms per dose; 200 micrograms per dose.
epinephrine (adrenaline)	Injection: 1 mg (as hydrochloride or hydrogen tartrate) in 1-mL ampoule.
<input type="checkbox"/> salbutamol	Injection: 50 micrograms (as sulfate)/mL in 5-mL ampoule. Metered dose inhaler (aerosol): 100 micrograms (as sulfate) per dose. Respirator solution for use in nebulizers: 5 mg (as sulfate)/mL.
26. SOLUTIONS CORRECTING WATER, ELECTROLYTE AND ACID-BASE DISTURBANCES	
26.1 Oral	
oral rehydration salts	See section 17.5.1.
potassium chloride	Powder for solution.

26.2 Parenteral	
glucose	Injectable solution: 5% (isotonic); 10% (hypertonic); 50% (hypertonic).
glucose with sodium chloride	Injectable solution: 5% glucose, 0.9% sodium chloride (equivalent to Na ⁺ 150 mmol/L and Cl ⁻ 150 mmol/L); 5% glucose, 0.45% sodium chloride (equivalent to Na ⁺ 75 mmol/L and Cl ⁻ 75 mmol/L).
potassium chloride	Solution for dilution: 7.5% (equivalent to K ⁺ 1 mmol/mL and Cl ⁻ 1 mmol/mL); 15% (equivalent to K ⁺ 2 mmol/mL and Cl ⁻ 2 mmol/mL).
sodium chloride	Injectable solution: 0.9% isotonic (equivalent to Na ⁺ 154 mmol/L, Cl ⁻ 154 mmol/L).
sodium hydrogen carbonate	Injectable solution: 1.4% isotonic (equivalent to Na ⁺ 167 mmol/L, HCO ₃ ⁻ 167 mmol/L). Solution: 8.4% in 10-mL ampoule (equivalent to Na ⁺ 1000 mmol/L, HCO ₃ ⁻ 1000 mmol/L).
□ sodium lactate, compound solution	Injectable solution.
26.3 Miscellaneous	
water for injection	2-mL; 5-mL; 10-mL ampoules.
27. VITAMINS AND MINERALS	
ascorbic acid	Tablet: 50 mg.
cholecalciferol*	Oral liquid: 400 IU/mL. Solid oral dosage form: 400 IU; 1000 IU. * Ergocalciferol can be used as an alternative.
iodine	Capsule: 200 mg. Iodized oil: 1 mL (480 mg iodine); 0.5 mL (240 mg iodine) in ampoule (oral or injectable); 0.57 mL (308 mg iodine) in dispenser bottle.
pyridoxine	Tablet: 25 mg (hydrochloride).
retinol	Capsule: 100 000 IU; 200 000 IU (as palmitate). Oral oily solution: 100 000 IU (as palmitate)/mL in multidose dispenser. Tablet (sugar-coated): 10 000 IU (as palmitate). Water-miscible injection: 100 000 IU (as palmitate) in 2-mL ampoule.
riboflavin	Tablet: 5 mg.
sodium fluoride	In any appropriate topical formulation.
thiamine	Tablet: 50 mg (hydrochloride).
<i>Complementary List</i>	
<i>calcium gluconate</i>	Injection: 100 mg/mL in 10-mL ampoule.
28. EAR, NOSE AND THROAT MEDICINES	

acetic acid	Topical: 2%, in alcohol.
<input type="checkbox"/> budesonide	Nasal spray: 100 micrograms per dose.
<input type="checkbox"/> ciprofloxacin	Topical: 0.3% drops (as hydrochloride).
<input type="checkbox"/> xylometazoline ^a	Nasal spray: 0.05%. ^a Not in children less than 3 months.
29. SPECIFIC MEDICINES FOR NEONATAL CARE	
29.1 Medicines administered to the neonate	
caffeine citrate	Injection: 20 mg/mL (equivalent to 10 mg caffeine base/mL). Oral liquid: 20 mg/mL (equivalent to 10 mg caffeine base/mL).
chlorhexidine	Solution or gel: 7.1% (digluconate) delivering 4% chlorhexidine (for umbilical cord care).
<i>Complementary List</i>	
<input type="checkbox"/> ibuprofen	Solution for injection: 5 mg/mL.
<input type="checkbox"/> prostaglandin E	Solution for injection: <i>Prostaglandin E1:</i> 0.5 mg/mL in alcohol. <i>Prostaglandin E2:</i> 1 mg/mL.
surfactant	Suspension for intratracheal instillation: 25 mg/mL or 80 mg/mL.
30. MEDICINES FOR DISEASES OF JOINTS	
30.1 Medicines used to treat gout	
30.2 Disease-modifying agents used in rheumatoid disorders (DMARDs)	
<i>Complementary List</i>	
hydroxychloroquine	Solid oral dosage form: 200 mg (as sulfate).
methotrexate	Tablet: 2.5 mg (as sodium salt).
30.3 Juvenile joint diseases	
acetylsalicylic acid* (acute or chronic use)	Suppository: 50 mg to 150 mg. Tablet: 100 mg to 500 mg. * For use for rheumatic fever, juvenile arthritis, Kawasaki disease.

Table 1.1: Medicines with age and weight restrictions

atazanavir	>25 kg
atropine	>3 months
benzyl benzoate	>2 years
betamethasone topical preparations	Hydrocortisone preferred in neonates
cefazolin	>1 month
ceftriaxone	>41 weeks corrected gestational age
darunavir	> 3 years
diloxanide	>25 kg
doxycycline	>8 years (except for serious infections e.g. cholera)
efavirenz	>3 years or >10 kg
emtricitabine	>3 months
fluoxetine	> 8 years
ibuprofen	>3 months (except IV form for patent <i>ductus arteriosus</i>)
mefloquine	>5 kg or >3 months
metoclopramide	Not in neonates
nevirapine	> 6 weeks
ondansetron	>1 month
propofol	>1 month
saquinavir	>25 kg
silver sulfadiazine	>2 months
tetracaine	Not in preterm neonates
trimethoprim	>6 months
xylometazoline	>3 months

Table 1.2: Explanation of dosage forms

A. Principal dosage forms used in EMLc - Oral administration

Term	Definition
Solid oral dosage form	<p>Refers to tablets or capsules or other solid dosage forms such as 'melts' that are immediate-release preparations. It implies that there is no difference in clinical efficacy or safety between the available dosage forms, and countries should therefore choose the form(s) to be listed depending on quality and availability.</p> <p>The term 'solid oral dosage form' is <i>never</i> intended to allow any type of modified-release tablet.</p>
Tablet	<p>Refers to:</p> <ul style="list-style-type: none"> • uncoated or coated (film-coated or sugar-coated) tablets that are intended to be swallowed whole; • unscored and scored*; • tablets that are intended to be chewed before being swallowed; • tablets that are intended to be dispersed or dissolved in water or another suitable liquid before being swallowed; • tablets that are intended to be crushed before being swallowed. <p>The term 'tablet' without qualification is <i>never</i> intended to allow any type of modified-release tablet.</p>
Tablet (qualified)	<p>Refers to a specific type of tablet:</p> <p>chewable - tablets that are intended to be chewed before being swallowed;</p> <p>dispersible - tablets that are intended to be dispersed in water or another suitable liquid before being swallowed;</p> <p>soluble - tablets that are intended to be dissolved in water or another suitable liquid before being swallowed;</p> <p>crushable - tablets that are intended to be crushed before being swallowed;</p> <p>scored - tablets bearing a break mark or marks where sub-division is intended in order to provide doses of less than one tablet;</p> <p>sublingual - tablets that are intended to be placed beneath the tongue.</p> <p>The term 'tablet' is <i>always</i> qualified with an additional term (in parentheses) in entries where one of the following types of tablet is intended: gastro-resistant (such tablets may sometimes be described as enteric-coated or as delayed-release), prolonged-release or another modified-release form.</p>

* Scored tablets may be divided for ease of swallowing, provided dose is a whole number of tablets.

Term	Definition
Capsule	Refers to hard or soft capsules. The term 'capsule' without qualification is <i>never</i> intended to allow any type of modified-release capsule.
Capsule (qualified)	The term 'capsule' with qualification refers to gastro-resistant (such capsules may sometimes be described as enteric-coated or as delayed-release), prolonged-release or another modified-release form.
Granules	Preparations that are issued to patient as granules to be swallowed without further preparation, to be chewed, or to be taken in or with water or another suitable liquid. The term 'granules' without further qualification is <i>never</i> intended to allow any type of modified-release granules.
Oral powder	Preparations that are issued to patient as powder (usually as single-dose) to be taken in or with water or another suitable liquid.
Oral liquid	Liquid preparations intended to be <i>swallowed</i> i.e. oral solutions, suspensions, emulsions and oral drops, including those constituted from powders or granules, but <i>not</i> those preparations intended for <i>oromucosal administration</i> e.g. gargles and mouthwashes. Oral liquids presented as powders or granules may offer benefits in the form of better stability and lower transport costs. If more than one type of oral liquid is available on the same market (e.g. solution, suspension, granules for reconstitution), they may be interchanged and in such cases should be bioequivalent. It is preferable that oral liquids do not contain sugar and that solutions for children do not contain alcohol.

B. Principal dosage forms used in EMLc - Parenteral administration

Term	Definition
Injection	Refers to solutions, suspensions and emulsions including those constituted from powders or concentrated solutions.
Injection (qualified)	Route of administration is indicated in parentheses where relevant.
Injection (oily)	The term 'injection' is qualified by '(oily)' in relevant entries.
Intravenous infusion	Refers to solutions and emulsions including those constituted from powders or concentrated solutions.

C. Other dosage forms

Mode of administration	Term to be used
To the eye	Eye drops, eye ointments.
Topical	For liquids: lotions, paints. For semi-solids: cream, ointment.
Rectal	Suppositories, gel or solution.
Vaginal	Pessaries or vaginal tablets.
Inhalation	Powder for inhalation, pressurized inhalation, nebulizer.

INDEX

abacavir (ABC).....	11	chloramphenicol.....	8
abacavir + lamivudine	12	chlorhexidine	22
acetic acid	29	chlorine base compound	22
acetylcysteine.....	5	chloroquine	14
<i>acetylsalicylic acid</i>	30	chloroxylenol	22
aciclovir	11, 26	<i>chlorpromazine</i>	27
albendazole	6, 7	cholecalciferol	29
alcohol based hand rub	22	cholera vaccine	26
<i>amikacin</i>	10	<i>ciclosporin</i>	16
amitriptyline	4	ciprofloxacin	9, 29
amodiaquine	13	<i>cisplatin</i>	16
amoxicillin.....	7	<i>clindamycin</i>	9
amoxicillin + clavulanic acid.....	7	clofazimine.....	9
amphotericin B	11, 13	cloxacillin	8
ampicillin.....	7	<i>coagulation factor IX</i>	20
antitetanus immunoglobulin	20	<i>coagulation factor VIII</i>	20
antivenom immunoglobulin.....	25	coal tar	21
artemether	13	cyclizine.....	4
artemether + lumefantrine	13	<i>cyclophosphamide</i>	16
artesunate	13	<i>cycloserine</i>	10
artesunate + amodiaquine	14	<i>cytarabine</i>	17
artesunate + mefloquine	14	<i>dacarbazine</i>	17
ascorbic acid.....	29	<i>dactinomycin</i>	17
<i>asparaginase</i>	16	dapsone	9
atazanavir.....	12	darunavir.....	12
atropine.....	3, 5, 27	<i>deferoxamine</i>	5, 19
<i>azathioprine</i>	16	<i>desmopressin</i>	19
azithromycin.....	8	dexamethasone.....	4, 5, 18, 23
<i>barium sulfate</i>	22	dextran 70.....	20
BCG vaccine.....	25	diazepam.....	4, 6
benzathine benzylpenicillin.....	7	diethylcarbamazine	7
benznidazole.....	15	digoxin.....	20
benzoyl peroxide.....	21	diloxanide.....	13
benzyl benzoate.....	21	<i>dimercaprol</i>	5
benzylpenicillin	7	diphtheria antitoxin	25
betamethasone	21	diphtheria vaccine.....	25
<i>bleomycin</i>	16	docusate sodium	4
budesonide.....	28, 29	<i>dopamine</i>	21
bupivacaine	3	<i>doxorubicin</i>	17
caffeine citrate.....	29	doxycycline	9, 14
calamine.....	21	efavirenz (EFV or EFZ).....	12
<i>calcium folinate</i>	16	eflornithine.....	15
calcium gluconate	5, 29	enalapril.....	20
<i>capreomycin</i>	10	entecavir	13
carbamazepine.....	6	epinephrine (adrenaline).....	5, 27, 28
<i>carboplatin</i>	16	erythromycin	9
cefalexin.....	7	ethambutol	10
cefazolin.....	7	ethanol	22
<i>cefotaxime</i>	8	<i>ethionamide</i>	10
<i>ceftazidime</i>	8	<i>ethosuximide</i>	6
ceftriaxone	7	<i>etoposide</i>	17
charcoal, activated.....	5	ferrous salt.....	19

<i>filgrastim</i>	17	<i>linezolid</i>	10
fluconazole.....	11	lopinavir + ritonavir (LPV/r).....	12
flucytosine.....	11	loratadine.....	5
fludrocortisone.....	24	lorazepam.....	6
fluorescein.....	21	<i>Lugol's solution</i>	24
fluoxetine.....	4, 27	<i>mannitol</i>	22
folic acid.....	19	measles vaccine.....	25
<i>fomepizole</i>	5	mebendazole.....	6
fresh–frozen plasma.....	19	mefloquine.....	14
furosemide.....	20, 22	<i>melarsoprol</i>	15
gentamicin.....	9, 26	meningococcal meningitis vaccine.....	26
glucagon.....	24	<i>mercaptapurine</i>	17
glucose.....	28	<i>mesna</i>	18
glucose with sodium chloride.....	28	<i>metformin</i>	24
glutaral.....	22	<i>methotrexate</i>	18, 30
griseofulvin.....	11	<i>methylprednisolone</i>	18
Haemophilus influenzae type b vaccine.....	25	metoclopramide.....	23
<i>haloperidol</i>	27	metronidazole.....	9, 13
halothane.....	3	miconazole.....	21
<i>heparin sodium</i>	19	midazolam.....	3, 4, 6
hepatitis A vaccine.....	26	miltefosine.....	13
hepatitis B vaccine.....	25	morphine.....	3, 4
HPV vaccine.....	25	mumps vaccine.....	26
<i>hydrochlorothiazide</i>	22	mupirocin.....	21
hydrocortisone.....	5, 18, 21, 24	naloxone.....	5
hydroxocobalamin.....	19	neostigmine.....	26
<i>hydroxycarbamide</i>	19	nevirapine (NVP).....	12
<i>hydroxychloroquine</i>	30	niclosamide.....	7
hyoscine hydrobromide.....	4	nifurtimox.....	15
ibuprofen.....	3, 15, 30	nitrofurantoin.....	9
<i>ifosfamide</i>	17	nitrous oxide.....	3
<i>imipenem + cilastatin</i>	8	<i>normal immunoglobulin</i>	20
influenza vaccine.....	26	nystatin.....	11
insulin injection (soluble).....	24	ofloxacin.....	27
intermediate-acting insulin.....	24	omeprazole.....	22
<i>intraperitoneal dialysis solution (of appropriate composition)</i>	27	ondansetron.....	4, 23
iodine.....	29	oral rehydration salts.....	23, 28
isoflurane.....	3	oseltamivir.....	12
isoniazid.....	10	<i>oxamniquine</i>	7
ivermectin.....	7	oxygen.....	3
Japanese encephalitis vaccine.....	26	packed red blood cells.....	19
<i>kanamycin</i>	10	<i>p-aminosalicylic acid</i>	10
ketamine.....	3	<i>pancreatic enzymes</i>	22
lactulose.....	4	paracetamol.....	3, 15
lamivudine (3TC).....	11	paromomycin.....	13
lamivudine + nevirapine + stavudine.....	12	pentamidine.....	15
lamivudine + nevirapine + zidovudine.....	12	permethrin.....	21
lamivudine + zidovudine.....	12	pertussis vaccine.....	25
levamisole.....	6	phenobarbital.....	6
<i>levofloxacin</i>	10	phenoxymethylpenicillin.....	8
levothyroxine.....	24	phenytoin.....	6
lidocaine.....	3	phytomenadione.....	19
lidocaine + epinephrine (adrenaline).....	3	platelet concentrates.....	19
		pneumococcal vaccine.....	25

podophyllum resin.....	21	sodium hydrogen carbonate.....	28
poliomyelitis vaccine	25	sodium lactate, compound solution	28
potassium chloride.....	28	sodium stibogluconate or meglumine	
<i>potassium iodide</i>	11, 24	antimoniate	13
potassium permanganate.....	21	<i>spironolactone</i>	22
povidone iodine.....	22	stavudine (d4T)	11
praziquantel.....	7	<i>streptomycin</i>	10
prednisolone	5, 18, 27	<i>succimer</i>	5
primaquine.....	14	sulfadiazine.....	15
procaine benzylpenicillin.....	8	sulfadoxine + pyrimethamine.....	14
proguanil.....	14	sulfamethoxazole + trimethoprim.....	9, 15
propofol.....	3	suramin sodium	15
propranolol	16	<i>surfactant</i>	30
<i>propylthiouracil</i>	24	suxamethonium.....	26
<i>prostaglandin E</i>	30	terbinafine	21
<i>protamine sulfate</i>	19	tetanus vaccine	25
pyrantel.....	7	tetracaine	27
pyrazinamide.....	10	tetracycline.....	27
<i>pyridostigmine</i>	26	thiamine.....	29
pyridoxine	29	<i>thioguanine</i>	18
pyrimethamine	15	tick-borne encephalitis vaccine	26
quinine.....	14	triclabendazole	7
rabies immunoglobulin.....	20	trimethoprim	9
rabies vaccine.....	26	tropicamide.....	22
ranitidine	23	tuberculin, purified protein derivative (PPD).....	24
retinol.....	29	typhoid vaccine	26
ribavirin	12	urea	21
riboflavin	29	valproic acid (sodium valproate).....	6
rifampicin.....	9, 10	<i>vancomycin</i>	9
rifapentine	10	varicella vaccine	26
ritonavir.....	12	vecuronium.....	26
rotavirus vaccine	25	<i>vinblastine</i>	18
rubella vaccine	25	<i>vincristine</i>	18
salbutamol.....	28	<i>warfarin</i>	19
salicylic acid	21	water for injection	29
senna	5	whole blood	20
silver sulfadiazine	21	xylometazoline	29
<i>sodium calcium edetate</i>	5	yellow fever vaccine	26
sodium chloride.....	28	zidovudine (ZDV or AZT).....	11
sodium fluoride.....	29	zinc sulfate	23